

MesenCult™-ACF Plus Medium

Animal component-free medium for human mesenchymal stem cells

Catalog #05445 1 Kit
#05448 1 Kit



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

MesenCult™-ACF Plus Medium is a standardized, animal component-free (ACF) and serum-free medium for the isolation and culture of human mesenchymal stromal cells, also known as mesenchymal stem cells (MSCs) from bone marrow (BM-MSCs) and adipose tissue (ADSCs). MesenCult™-ACF Plus Medium is optimized for the expansion of MSCs in vitro as well as their enumeration using the colony-forming unit-fibroblast (CFU-F) assay. MesenCult™-ACF Plus Medium supports the isolation and long-term growth of human bone marrow- and adipose tissue-derived MSCs, and cells maintain robust multi-lineage differentiation potential in vitro.

MesenCult™-ACF Plus Medium also supports the expansion of human ES- or iPS-derived mesenchymal progenitor cells (MPCs), and is included in STEMdiff™ Mesenchymal Progenitor Kit (Catalog #05240).

MesenCult™-ACF Plus Medium must be used in conjunction with Animal Component-Free Cell Attachment Substrate (Component #07130) and Animal Component-Free Cell Dissociation Kit (Catalog #05426), providing a complete, defined ACF culture system. Components of Animal Component-Free Cell Attachment Substrate and Animal Component-Free Cell Dissociation Kit are pre-screened and tested for optimal cell adherence when cells are cultured with MesenCult™-ACF Plus Medium.

For animal component-free and optimized cryopreservation, MesenCult™-ACF Freezing Medium (Catalog #05490) is recommended for human MSCs previously cultured in MesenCult™ media, including MesenCult™-ACF Plus. For a complete list of related products, including differentiation media available, visit www.stemcell.com, or contact us at techsupport@stemcell.com.

NOTE: Complete MesenCult™-ACF Plus Medium must be supplemented with L-Glutamine (Catalog #07100); see Preparation of Reagents and Materials.

Ordering Information

PRODUCT NAME	CATALOG #	SIZE	COMPONENTS
MesenCult™-ACF Plus Medium Kit	05445	1 Kit	<ul style="list-style-type: none">MesenCult™-ACF Plus MediumMesenCult™-ACF Plus 500X Supplement
MesenCult™-ACF Plus Culture Kit	05448	1 Kit	<ul style="list-style-type: none">MesenCult™-ACF Plus MediumMesenCult™-ACF Plus 500X SupplementAnimal Component-Free Cell Attachment Substrate

Components

The following components are available as part of a kit (Catalog #05445 or #05448) and are not available for individual sale.

COMPONENT NAME	COMPONENT #	SIZE	STORAGE	SHELF LIFE
MesenCult™-ACF Plus Medium	05446	500 mL	Store at 2 - 8°C.	Stable for 12 months from date of manufacture (MFG) on label.
MesenCult™-ACF Plus 500X Supplement	05447	1 mL	Store at -20°C.	Stable for 2 years from date of manufacture (MFG) on label.
Animal Component-Free Cell Attachment Substrate	07130	1 mL	Store at 2 - 8°C.	Stable until expiry date (EXP) on label.

None of the above components contain antibiotics.

Materials Required but Not Included

PRODUCT NAME	CATALOG #
0.5 mL screw cap polypropylene tubes	e.g. Sarstedt 72.785.005
3% Acetic Acid with Methylene Blue	07060
Animal Component-Free Cell Dissociation Kit <ul style="list-style-type: none"> • ACF Enzymatic Dissociation Solution • ACF Enzyme Inhibition Solution 	05426
Collagenase Type I (0.25%) OR Collagenase A, ACF	07902 OR 07434
D-PBS (Without Ca++ and Mg++)	37350
Hausser Scientific™ Bright-Line Hemocytometer	100-1181
L-Glutamine	07100
Lymphoprep™	07801
Non-treated culture dish, 100 mm	e.g. 38045
Polypropylene conical tubes, 15 mL and 50 mL	e.g. 38009 and 38010
Tissue culture-treated 6-well plates	e.g. 38015
Trypan Blue	07050

Preparation of Reagents and Materials

Complete MesenCult™-ACF Plus Medium

Use sterile technique to prepare complete MesenCult™-ACF Plus Medium (MesenCult™-ACF Plus Medium + MesenCult™-ACF Plus 500X Supplement + L-Glutamine). The following example is for preparing 500 mL of complete medium. If preparing other volumes, adjust accordingly.

1. Thaw MesenCult™-ACF Plus 500X Supplement on ice for 1 - 2 hours or overnight at 2 - 8°C. Mix thoroughly.
NOTE: Once thawed, use immediately or aliquot and store at -20°C. For aliquoting, use 0.5 mL screw cap polypropylene tubes. Do not exceed the shelf life of the supplement. After thawing the aliquots, use immediately. **Do not re-freeze.**
2. Add 1 mL of MesenCult™-ACF Plus 500X Supplement to 500 mL of MesenCult™-ACF Plus Medium. Mix thoroughly.
3. Add L-Glutamine to a final concentration of 2 mM. Mix thoroughly.

NOTE: If not used immediately, store complete MesenCult™-ACF Plus Medium **at 2 - 8°C for up to 1 week**. Do not exceed the shelf life of the individual components.

Coating Cultureware with Animal Component-Free Cell Attachment Substrate

Use sterile technique when coating cultureware with Animal Component-Free Cell Attachment Substrate.

NOTE: Use only tissue culture-treated cultureware.

1. Dilute Animal Component-Free Cell Attachment Substrate in D-PBS (Without Ca++ and Mg++) as follows:
 - For MSCs after isolation, dilute Attachment Substrate 1 in 300 in D-PBS.
For example, add 20 µL of Substrate to 5.98 mL of D-PBS.
 - For isolation of MSCs from bone marrow, dilute Attachment Substrate 1 in 150 in D-PBS.
For example, add 40 µL of Substrate to 5.96 mL of D-PBS.
 - For isolation of MSCs from adipose tissue, dilute Attachment Substrate 1 in 300 in D-PBS.
For example, add 20 µL of Substrate to 5.98 mL of D-PBS.
2. Gently mix the diluted substrate solution. Do not vortex.
3. Immediately use the diluted substrate solution to coat cultureware. Refer to Table 1 for recommended coating volumes.

Table 1. Recommended Volumes for Coating Cultureware with Diluted Animal Component-Free Cell Attachment Substrate

CULTUREWARE	VOLUME OF DILUTED ANIMAL COMPONENT-FREE CELL ATTACHMENT SUBSTRATE SOLUTION
6-well plate	1 mL/well
T-25 cm ² flask	2.5 mL/flask
T-75 cm ² flask	6 mL/flask

- Gently tilt the cultureware to spread the substrate solution evenly across the surface.
- Incubate at room temperature (15 - 25°C) for at least 2 hours before use. Do not let the substrate solution evaporate.
NOTE: If not used immediately, the cultureware must be sealed to prevent evaporation of substrate solution (e.g. with Parafilm®). Sealed cultureware can be stored at 2 - 8°C for up to 3 days after coating. Allow stored coated cultureware to come to room temperature for 30 minutes before proceeding to step 6.
- Gently tilt the cultureware onto one side and allow excess substrate solution to collect at the edge. Remove the excess solution using a serological pipette or by aspiration. Ensure that the coated surface is not scratched.
- Wash cultureware once, using D-PBS (e.g. use 2 mL/well if using a 6-well plate).
- Aspirate wash solution when MSCs are ready to be plated.

Directions for Use

Please read the entire protocol before proceeding. Use sterile technique when performing the following protocols:

- Isolation of Human MSCs from Bone Marrow
- Isolation of Human MSCs from Adipose Tissue
- CFU-F Assay
- Expansion of Human MSCs

A) Isolation of Human MSCs from Bone Marrow (BM)

The following protocol is for isolating MSCs from 25 mL of freshly isolated human BM using density gradient medium (Lymphoprep™) separation. If using other volumes, adjust accordingly.

- Count nucleated cells in the BM sample using 3% Acetic Acid with Methylene Blue.
- Split the BM sample into 2 x 50 mL conical tubes (i.e. 12.5 mL of BM sample per tube).
- Add 22.5 mL of room temperature (15 - 25°C) D-PBS containing 2 mM EDTA per tube.
- To each of 3 x 50 mL new conical tubes, add 17 mL of Lymphoprep™.
- Layer 23 mL of the BM suspension (from step 3) on top of the Lymphoprep™ in each tube.
- Centrifuge tubes at 300 x g for 30 minutes with the **brake off**.
- Collect the mononuclear cell (MNC) layer at the plasma:Lymphoprep™ interface and place in a single new 50 mL conical tube.
NOTE: Sometimes it is difficult to see the cells at the interface. In this case, remove some of the Lymphoprep™ along with the enriched cells in order to maximize cell recovery.
- Wash cells with cold (2 - 8°C) D-PBS containing 2 mM EDTA.
- Centrifuge the tube at 300 x g for 10 minutes with the **brake on**.
- Discard supernatant and resuspend the cell pellet in complete MesenCult™-ACF Plus Medium (e.g. 2 - 4 mL of complete medium).
- Count nucleated cells using 3% Acetic Acid with Methylene Blue.

B) Isolation of Human MSCs from Adipose Tissue

- Add 2 - 4 mL of Collagenase Type I (0.25%) to the adipose tissue in a 100 mm dish.
NOTE: To maintain an ACF workflow, use Collagenase A, ACF, prepared at 0.2% in D-PBS containing 0.2% recombinant albumin.
- Finely mince tissue with a scalpel. Transfer minced tissue to a 50 mL conical tube.
- Add 5 mL of collagenase per cm³ of tissue. Incubate in a shaking water bath or shaking incubator at 37°C for 1 hour.
For example, use 15 mL of collagenase for 3 cm³ of tissue.
- Remove tube from the water bath or incubator. Place upright for 5 minutes to allow separation of the lipid layer from the aqueous layer.
- Using a pipettor or aspirator, remove and discard the top lipid layer.
- Add D-PBS containing 1 mM EDTA and 0.2% recombinant albumin (or 2% fetal bovine serum) to reach a final volume of 50 mL.
- Centrifuge cells at 300 x g for 10 minutes with the **brake on**.

8. Discard supernatant and resuspend the cell pellet in complete MesenCult™-ACF Plus Medium.
9. Count nucleated cells using 3% Acetic Acid with Methylene Blue.

C) CFU-F Assay

The following protocol is for setting up a CFU-F assay in a 6-well tissue culture-treated plate. If using other cultureware, adjust cell numbers and volumes accordingly.

NOTE: Only use tissue culture-treated cultureware.

1. Coat wells with Animal Component-Free Cell Attachment Substrate (see Preparation of Reagents and Materials).
2. Plate cells in 2 mL of complete MesenCult™-ACF Plus Medium per coated well. Plate cells at 3 - 4 different densities for each cell type used. Refer to Table 2 for recommended cell plating densities.

Table 2: Recommended Cell Plating Densities for Setting up the CFU-F Assay

CELL TYPE	PLATING DENSITY (cells/cm ²)	EXAMPLE OF NUMBER OF CELLS PER WELL OF A 6-WELL PLATE
Adipose-derived MSCs	0.5 - 4 x 10 ³	0.5 x 10 ⁴ 1 x 10 ⁴ 2 x 10 ⁴ 4 x 10 ⁴
BM-derived MSCs	1 - 4 x 10 ⁴	1 x 10 ⁵ 2 x 10 ⁵ 3 x 10 ⁵ 4 x 10 ⁵

3. Incubate at 37°C for 10 - 15 days until colonies (> 40 cells/colony) appear in the well.
4. Perform a half-medium change on day 7 (i.e. aspirate 1 mL of medium and add 1 mL of complete MesenCult™-ACF Plus Medium per well).
5. Fix, stain, and count the CFU-F colonies.

D) Expansion of Human MSCs

The following protocol is for culturing cells in a single T-25 cm² flask. If using other cultureware, adjust cell numbers and volumes accordingly.

NOTE: Only use tissue culture-treated cultureware.

1. Coat a T-25 cm² flask with Animal Component-Free Cell Attachment Substrate (see Preparation of Reagents and Materials).
2. Seed freshly isolated cells (prepared in section A or B) in 6 mL of complete MesenCult™-ACF Plus Medium per coated flask. Recommended densities are as follows:
 - For BM-derived cells: 4 - 10 x 10⁴ freshly isolated cells/cm² (i.e. 1 - 2.5 x 10⁶ cells per T-25 cm² flask)
 - For adipose-derived cells: 6 - 10 x 10³ freshly isolated cells/cm² (i.e. 1.5 - 2.5 x 10⁵ cells per T-25 cm² flask)
3. Incubate at 37°C until cells are approximately 80% confluent. This takes approximately 9 - 14 days.
NOTE: If needed, perform a half-medium change on day 7 (i.e. aspirate 3 mL of medium and add 3 mL of complete MesenCult™-ACF Plus Medium per flask).
4. Passage cells using Animal Component-Free Cell Dissociation Kit, as follows:
 - i. Coat a T-25 cm² flask with Animal Component-Free Cell Attachment Substrate (see Preparation of Reagents and Materials).
 - ii. Warm ACF Enzymatic Dissociation Solution and ACF Enzyme Inhibition Solution to room temperature (15 - 25°C). Do not incubate at 37°C.
 - iii. Wash cells once with 2.5 mL of D-PBS.
 - iv. Add 2.5 mL of ACF Enzymatic Dissociation Solution and incubate at 37°C for 3 - 6 minutes. Tap the flask to detach cells. If less than 90% of cells have detached, incubate at 37°C for an additional 1 - 2 minutes and tap the flask again.
 - v. Add 2.5 mL of ACF Enzyme Inhibition Solution and collect cells in a 15 mL conical polypropylene tube.
 - vi. Wash the flask with 5 mL of complete MesenCult™-ACF Plus Medium and place into the tube from step v.
 - vii. Centrifuge the tube at 300 x g for 8 minutes with the **brake on**.
 - viii. Discard the supernatant and resuspend the cell pellet in complete MesenCult™-ACF Plus Medium (see Notes and Tips).
 - ix. Count viable cells using Trypan Blue and a hemocytometer. Seed a coated T-25 cm² flask (prepared in step i) at a density of 1.5 - 4 x 10³ cells/cm² (i.e. 4 - 10 x 10⁴ cells/flask).

- x. Incubate at 37°C until the cells are approximately 80% confluent. This takes approximately 3 - 6 days.
5. Repeat step 4 as needed.

Notes and Tips

- The use of polypropylene tubes (e.g. Catalog #38009 and #38010) during subculture will help to prevent the MSCs from sticking to the tubes.
- To assess trilineage potential, MesenCult™ media are available for adipogenic differentiation (Catalog #05412), osteogenic differentiation (Catalog #05465), and chondrogenic differentiation (Catalog #05455).
- To break apart cell aggregates, use a 1 mL pipettor to gently pipette the cell pellet up and down a few times [section D, step 4 (viii)].

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED. FOR ADDITIONAL INFORMATION ON QUALITY AT STEMCELL, REFER TO WWW.STEMCELL.COM/COMPLIANCE.

Copyright © 2023 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and MesenCult are trademarks of STEMCELL Technologies Canada Inc. Parafilm is a registered trademark of Bemis Company, Inc. Lymphoprep is a trademark of Serumwerk Bernburg AG. The products sold under the Lymphoprep brand name are also manufactured by Serumwerk Bernburg AG. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.