

Anti-Human IL-4 Antibody, Clone MP4-25D2

Rat monoclonal antibody against human IL-4, unconjugated

Catalog #100-1454 100 μ g 2 mg/mL

Product Description

This rat monoclonal antibody (clone MP4-25D2) reacts with human interleukin 4 (IL-4). IL-4 is an ~15 kDa protein primarily composed of a four-helix bundle and is produced by a host of immune cells including T helper 2 (Th2) cells, basophils, eosinophils, and mast cells. It exhibits high affinity for the IL-4 receptor alpha (IL-4R α) chain, which dimerizes with either the common gamma chain (γ c) or IL-13 receptor alpha 1 (IL-13R α 1), leading to activation of downstream JAK-STAT signaling. IL-4 is involved in the differentiation of CD4+ T lymphocytes into Th2 cells, class switching of B cells to IgE, and activation of M2 macrophages with IL-13 via an alternative pathway. MP4-25D2 is a neutralizing antibody and is suitable for the detection of intracellular IL-4 by flow cytometry.

| Target Antigen: | IL-4 |
|---------------------|---|
| Alternative Names: | B cell stimulatory factor 1, BCDF, BCGF, BSF-1, interleukin-4 |
| Gene ID: | 3565 |
| Species Reactivity: | Human |
| Host Species: | Rat |
| Clonality: | Monoclonal |
| Clone: | MP4-25D2 |
| Isotype: | lgG1, kappa |
| Immunogen: | Purified recombinant human IL-4 |
| | |

Unconjugated

Conjugate:

Applications

Verified Applications: FC

Reported Applications: FA

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IHC-P: Immunohistochemistry (paraffin-embedded); IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Purification: The antibody was purified by affinity chromatography.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. Stable until expiry date (EXP) on label.

Directions for Use: For flow cytometry, the suggested use of this antibody is ≤ 0.25 μg per 1 x 10⁶ in 100 μL. It is

recommended that the antibody be titrated for optimal performance for each application.

Related Products

For a complete list of antibodies, including other conjugates, sizes, and clones, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/antibodies, or contact us at techsupport@stemcell.com.

References

De Fanis U et al. (2007) GATA3 up-regulation associated with surface expression of CD294/CRTH2: a unique feature of human Th cells. Blood 109(10): 4343-50.

Gadani SP et al. (2012) IL-4 in the brain: a cytokine to remember. J Immunol 189 (9): 4213-19.

Geha RS et al. (2003) The regulation of immunoglobulin E class-switch recombination. Nat Rev Immunol 3(9): 721-32.

Gordon S & Martinez FO. (2010) Alternative activation of macrophages: mechanism and functions. Immunity 32(5): 593-604.

Han K et al. (2021) Fasting-induced FOXO4 blunts human CD4+T helper cell responsiveness. Nat Metab 3(3): 318-26.

Heller NM et al. (2008) Type I IL-4 receptors selectively activate IRS-2 to induce target gene expression in macrophages. Sci Signal 1(51): ra17.

Li L et al. (2019) TLR8-mediated metabolic control of human Treg function: a mechanistic target for cancer immunotherapy. Cell Metab 29(1): 103–23.e5.

Nakatsukasa H et al. (2015) The DNA-binding inhibitor Id3 regulates IL-9 production in CD4+ T cells. Nat Immunol 16(10): 1077-84.

Powers R et al. (1992) Three-dimensional solution structure of human interleukin-4 by multidimensional heteronuclear magnetic resonance spectroscopy. Science 256(5064): 1673–77.

Silva-Filho JL et al. (2014) IL-4: an important cytokine in determining the fate of T cells. Biophys Rev 6(1): 111-18.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2024 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.