

Anti-Human Perforin Antibody, Clone deltaG9

Mouse monoclonal antibody against human perforin, unconjugated

Catalog #100-1457

100 µg

1.0 mg/mL

Product Description

This mouse monoclonal antibody (clone deltaG9) reacts with human perforin. Perforin is an ~70 kDa protein consisting of four domains: the N- and C-termini, which are related to its function, and two central domains that share similarities to complement proteins C6 - C9. Perforin is involved in the perforin/granzyme apoptosis pathway and is produced by natural killer and cytotoxic T cells. Upon contact with a target cell, perforin monomers are released and aggregate to form pores in the membrane of the target cell, disrupting its membrane integrity. This allows granzyme B to enter the target cell and initiate apoptosis of virally infected and transformed cells. The deltaG9 antibody is suitable for the detection of intracellular perforin by flow cytometry.

Target Antigen:	Perforin
Alternative Names:	Cytolysin, lymphocyte pore forming protein, P1, perforin 1, PFP
Gene ID:	5551
Species Reactivity:	Human
Host Species:	Mouse
Clonality:	Monoclonal
Clone:	deltaG9
Isotype:	lgG2b, kappa
Immunogen:	Purified granules from human YT lymphoma cell line
Conjugate:	Unconjugated

Applications

Reported Applications: FC, IF, IHC

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IHC-P: Immunohistochemistry (paraffin-embedded); IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Purification:	The antibody was purified by affinity chromatography.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Stable until expiry date (EXP) on label.
Directions for Use:	For flow cytometry, the suggested use of this antibody is 0.08 μ g per 5 x 10 ⁵ cells in 80 μ L. It is recommended that the antibody be titrated for optimal performance for each application.

Related Products

For a complete list of antibodies, including other conjugates, sizes, and clones, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/antibodies, or contact us at techsupport@stemcell.com.

References

Jacob T et al. (2001) Examination of the apoptotic pathway and proteolysis in the pathogenesis of popliteal artery aneurysms. Eur J Vasc Endovasc Surg 22(1): 77–85.

Liu CC et al. (1995). Perforin: structure and function. Immunol Today 16(4): 194-201.

Liu X et al. (2022) Single-cell transcriptomics links malignant T cells to the tumor immune landscape in cutaneous T cell lymphoma. Nat Commun 13(1): 1158.

Lund H et al. (2013) Natural killer cells in afferent lymph express an activated phenotype and readily produce IFN-y. Front Immunol 4: 395.

Narazaki H et al. (2003) Perforin-dependent killing of tumor cells by V γ 1V δ 1-bearing T-cells. Immunol Lett 86(1): 113–9.

Osińska I et al. (2014) Perforin: an important player in immune response. Cent Eur J Immunol 39(1): 109-15.

Trapani JA & Smyth MJ (2002) Functional significance of the perforin/granzyme cell death pathway. Nat Rev Immunol 2(10): 735-47.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2024 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.