

# StemSpan™ Myeloid Expansion Supplement II (100X)



Scientists Helping Scientists™ | [WWW.STEMCELL.COM](http://WWW.STEMCELL.COM)

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

[INFO@STEMCELL.COM](mailto:INFO@STEMCELL.COM) • [TECHSUPPORT@STEMCELL.COM](mailto:TECHSUPPORT@STEMCELL.COM)

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog #02694 1 mL

## Product Description

StemSpan™ Myeloid Expansion Supplement II (100X) contains a combination of recombinant human cytokines formulated to selectively promote the expansion and differentiation of human myeloid progenitor cells, generating large numbers of CD14+ monocytes from CD34+ cells isolated from human cord blood (CB) samples.

StemSpan™ Myeloid Expansion Supplement II (100X) is intended for use in combination with any of the following StemSpan™ media:

- StemSpan™ SFEM (Catalog #09600)
- StemSpan™ SFEM II (Catalog #09605)
- StemSpan™-XF (Catalog #100-0073)
- StemSpan™-AOF (Catalog #100-0130)

### Advantages:

- Promotes the production of hundreds of CD14+ monocytes per input CD34+ human CB cell in 14-day liquid cultures.
- Supplied as a 100X concentrate. After thawing and mixing, the tube contents can be added directly to any hematopoietic cell expansion medium.

## Properties

**Storage:** Store at -20°C.

**Shelf Life:** Stable for 15 months from date of manufacture (MFG) on label.

- Contains:**
- Recombinant human fms-like tyrosine kinase 3 ligand (Flt3L)
  - Recombinant human stem cell factor (SCF)
  - Recombinant human thrombopoietin (TPO)
  - Recombinant human macrophage colony-stimulating factor (M-CSF)
  - Recombinant human granulocyte/macrophage colony-stimulating factor (GM-CSF)
  - Supplements

## Directions for Use

### PREPARATION OF COMPLETE MEDIUM

1. Thaw StemSpan™ Myeloid Expansion Supplement II (100X) at room temperature (15 - 25°C) until just thawed. Mix thoroughly.

NOTE: If necessary, centrifuge for 30 seconds to recover liquid from cap.

NOTE: Once thawed, store supplement at 2 - 8°C for up to 1 month. Alternatively, aliquot and store at -20°C. After thawing aliquots, do not re-freeze.

2. Add StemSpan™ Myeloid Expansion Supplement II (100X) to culture medium at a 1 in 100 dilution (e.g. add 1 mL of Supplement to 99 mL of culture medium). Mix thoroughly.

### RECOMMENDED PROTOCOL FOR MONOCYTE EXPANSION WITH StemSpan™ MEDIA

For optimal performance, use one of the following StemSpan™ media in this protocol:

- StemSpan™ SFEM II (Catalog #09605)

1. Thaw StemSpan™ medium at room temperature (15 - 25°C) or overnight at 2 - 8°C. Mix thoroughly.

NOTE: If not used immediately, aliquot into tubes and store at -20°C. Once aliquots are thawed, do not re-freeze.

2. Prepare complete medium as described in the Preparation section, using a StemSpan™ medium as the culture medium.

3. Thaw cryopreserved CD34+ cells, or use an EasySep™ kit to isolate CD34+ cells from fresh whole CB or from cryopreserved CB mononuclear cells (MNCs), as indicated below:
  - For fresh whole CB, use EasySep™ Human Cord Blood CD34 Positive Selection Kit II (Catalog #17896)
  - For frozen CB MNCs, use EasySep™ Human CD34 Positive Selection Kit II (Catalog #17856)
4. **Day 0:** Plate CD34+ cells in complete medium at  $1 \times 10^4$  cells/mL. Refer to Table 1 for recommended plating concentrations; optimal cell concentrations and cultureware are dependent on experimental objectives and cell quality.

**Table 1. Recommended CD34+ Cell Concentrations for Various Cultureware**

CULTUREWARE*	VOLUME OF MEDIUM PER WELL	NUMBER OF CELLS PER WELL
6-well plate	2 mL	$2 \times 10^4$
24-well plate	1 mL	$1 \times 10^4$
96-well plate	100 $\mu$ L	$1 \times 10^3$

\*Both tissue culture-treated and non-tissue culture-treated are suitable.

5. Incubate at 37°C and 5% CO<sub>2</sub>.
6. **Day 3 or 4:** Add an equal volume of fresh complete medium (i.e. if using a 24-well plate, add 1 mL complete medium per well).
7. **Day 7:** Harvest and replate in fresh complete medium at  $< 1 \times 10^5$  cells/mL. Alternatively, add fresh complete medium to maintain cell concentration at  $< 1 \times 10^5$  cells/mL.
8. **Day 10/11:** Add an equal volume of fresh complete medium (i.e. if using a 24-well plate, add 1 mL complete medium per well) to maintain cell concentration at  $< 1 \times 10^5$  cells/mL.
9. **Day 14:** Harvest cells for evaluation or downstream applications. Count total viable cells using Trypan Blue (Catalog #07050) and a hemocytometer (or an automated cell counting method), and measure monocyte progenitor cell surface marker expression by flow cytometry.

NOTE: Cultures can be continued beyond 14 days with periodic dilution every 3 - 4 days to maintain a cell concentration of  $1 \times 10^5$  cells/mL.

## Notes and Tips

### RELATED PRODUCTS

For related products, including specialized culture and storage media, supplements, antibodies, cytokines, and small molecules, visit [www.stemcell.com/HSPCworkflow](http://www.stemcell.com/HSPCworkflow) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com). For available fresh and cryopreserved peripheral blood, cord blood, and bone marrow products in your region, visit [www.stemcell.com/primarycells](http://www.stemcell.com/primarycells).

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED. FOR ADDITIONAL INFORMATION ON QUALITY AT STEMCELL, REFER TO [WWW.STEMCELL.COM/COMPLIANCE](http://WWW.STEMCELL.COM/COMPLIANCE).

Copyright © 2021 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, EasySep, and StemSpan are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.