

Dissociation Reagents

Collagenase Type V

For digestion of native collagen fibrils



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog #	07430	100 mg
	07431	1 g
	100-0681	5 g

Product Description

Collagenase is a protease consisting of a single polypeptide chain approximately 1000 amino acid residues in length. Collagenase is capable of digesting native collagen fibrils commonly found in connective tissues and therefore is frequently used for tissue dissociation. Collagenase preparations contain the activity of several proteases, including collagenase, caseinase, clostripain, and trypsin (Kessler & Yaron). Collagenase Type V contains high levels of collagenase and caseinase activities.

Product Information

Alternative Names:	Clostridium histolyticum collagenase; Collagenase 5; Collagenase Type 5; Collagenase V
Format:	Lyophilized powder
Storage:	Store at 2 - 8°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Reconstitution:	Dissociation reagents can be reconstituted in a balanced salt solution or buffer of choice.
Molecular Weight:	68 - 130 kDa
CAS Number:	9001-12-1
Optimum pH:	6.3 - 8.5
Cleavage Site:	-Pro-X-†-Gly-Pro-Y- : X = neutral Y = nonspecific

Please refer to the Safety Data Sheet (SDS) for hazard information.

Specifications

Source:	Clostridium histolyticum
Activity:	Collagenase: ≥ 450 CDU/mg dry weight (mgdw); Caseinase: ≥ 450 u/mgdw; Clostripain: ≤ 4.0 u/mgdw; Trypsin: ≤ 0.3 u/mgdw. See Notes for further information.

Related Products

For a complete list of dissociation reagents, as well as related products available from STEMCELL Technologies, visit www.stemcell.com or contact us at techsupport@stemcell.com.

Notes

ACTIVITY UNITS

Collagenase: 1 collagenase digestion unit (CDU) equals 1 μ mol of L-leucine equivalents released from collagen in 5 hours at 37°C, pH 7.5.

Caseinase: 1 unit equals 1 μ mol of L-leucine equivalents released from 25 mg vitamin-free casein in 5 hours at 37°C, pH 7.5; measures non-specific proteolytic activity.

Clostripain: 1 unit hydrolyzes 1 μ mol of N α -benzoyl-L-arginine ethyl ester (BAEE)/minute at 25°C at pH 7.6, after activation in 2.5 mM dithiothreitol (DTT).

Trypsin: 1 unit hydrolyzes 1 μ mol of BAEE/minute at 25°C at pH 7.6.

References

Barrett AJ & Starkey PM. (1973) The interaction of α 2-macroglobulin with proteinases. J Biochem 133: 709–24.

Kessler E & Yaron A. (1973) A novel aminopeptidase from clostridium histolyticum. Biochem Biophys Res Commun 50(2): 405–12.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2021 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.