

Dissociation Reagents

Dispase, ACF

Animal component-free non-specific protease

Catalog # 07446
100-0396

10 mg
50 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Dispase, Animal Component-Free (ACF) is a neutral, amino-endoprotease that cleaves the N-terminal peptide bond of non-polar amino acid residues and is obtained from cultures free of animal-derived materials. Dispase has mild proteolytic activity, which makes it especially useful for the isolation and passaging of primary cells. Its proteolytic activity also allows it to maintain membrane integrity. Dispase is commonly used with other proteases such as collagenase in cell isolation and for dissociation of tissues such as neural (Dietrich et al.), kidney (Presnell et al.), epithelial (Smoot et al.), endothelial (Müller et al.), lung (Barkauskas et al.), and colon (Roig et al.), as well as dissociation of stem cells (Salmon et al.; Thomson et al.).

Product Information

Alternative Names:	Neutral protease; Proteinase
Format:	Lyophilized powder
Storage:	Store at 2 - 8°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Reconstitution:	Dissociation reagents can be reconstituted in a balanced salt solution or buffer of choice.
Molecular Weight:	32.5 kDa
CAS Number:	42613-33-2
Optimum pH:	5.9 - 7.0
Cleavage Site:	-X- † -Leu/Phe- † -Y : X/Y = nonspecific

Specifications

Source:	Bacillus polymyxa
Activity:	≥ 4 units/mg dry weight. See Notes for further information.

Related Products

For a complete list of dissociation reagents, as well as related products available from STEMCELL Technologies, visit www.stemcell.com or contact us at techsupport@stemcell.com.

Notes

ACTIVITY UNITS

1 unit releases 1 μmol folin-positive amino acids equivalent to 1 μmol tyrosine/minute from casein at 37°C, pH 7.5.

References

- Barkauskas CE et al. (2013) Type 2 alveolar cells are stem cells in adult lung. *J Clin Invest* 123(7): 3025–36.
- Dietrich J et al. (2005) EIF2B5 mutations compromise GFAP+ astrocyte generation in vanishing white matter leukodystrophy. *Nat Med* 11(3): 277–83.
- Müller AM et al. (2002) Expression of the endothelial markers PECAM-1, vWf, and CD34 in vivo and in vitro. *Exp Mol Pathol* 72(3): 221–9.
- Presnell SC et al. (2011) Isolation, characterization, and expansion methods for defined primary renal cell populations from rodent, canine, and human normal and diseased kidneys. *Tissue Eng C, Methods* 17(3): 261–73.
- Roig AI et al. (2010) Immortalized epithelial cells derived from human colon biopsies express stem cell markers and differentiate in vitro. *Gastroenterology* 138(3): 1012–21.e5.
- Salmon B et al. (2013) MEPE-derived ASARM peptide inhibits odontogenic differentiation of dental pulp stem cells and impairs mineralization in tooth models of X-linked hypophosphatemia. *PLoS One* 8(2): e56749.
- Smoot DT et al. (2000) A method for establishing primary cultures of human gastric epithelial cells. *Methods Cell Sci* 22(2-3): 133–6.
- Thomson JA. (1998) Embryonic stem cell lines derived from human blastocysts. *Science* 282(5391): 1145–7.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2020 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.