#### Anti-Human CD45 Antibody, Clone HI30, PE

## **Antibodies**

Mouse monoclonal IgG1 antibody against human, chimpanzee CD45,

PE-conjugated

Catalog #60018PE #60018PE.1

100 Tests 5 μL/test 25 Tests 5 μL/test



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

# **Product Description**

The HI30 antibody reacts with all isoforms of CD45, a type I transmembrane glycoprotein expressed on the surface of most hematopoietic cells except mature erythrocytes, platelets, and plasma cells; expression of CD45 is lost during differentiation of these cell types. CD45 is a member of the protein tyrosine phosphatase family and functions in a number of immunoregulatory processes, including cell activation, growth, differentiation, and oncogenic transformation. The large cytoplasmic portion of CD45 contains two tyrosine phosphatase domains (one of which is enzymatically active) that are involved in modulating the function of intracellular substrates such as the Src kinases Lck and Fyn. Several isoforms of CD45 have been identified, the expression of which differs according to cell type and functional status. Alternative splicing of three exons (4, 5, 6) encoding the extracellular RA, RB, and RC polypeptide sequences gives rise to up to 8 isoforms with molecular masses in the range of 180 - 240 kDa. The leukocyte common antigen (LCA), the region recognized by the HI30 antibody, is an extracellular region located proximal to the membrane and common to all isoforms of CD45.

Target Antigen Name: CD45

Alternative Names: B220, LCA, T200

Gene ID: 5788

Species Reactivity: Human, Chimpanzee

Host Species: Mouse
Clonality: Monoclonal
Clone: HI30

Isotype: IgG1, kappa

Immunogen: Full-length human CD45 protein

Conjugate: PE (Phycoerythrin)

# **Applications**

Verified: FC Reported: FC

Special Applications: This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including

EasySep™ Human T Cell Enrichment Kit (Catalog #19051), EasySep™ Human CD4+ T Cell Enrichment Kit (Catalog #19052) and EasySep™ Human CD45 Depletion Kit (Catalog #18259; partial blocking may be

observed).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

# **Properties**

Formulation: Phosphate-buffered saline, pH 7.2, containing 0.09% sodium azide, 0.1% gelatin, and < 0.2% (w/v) bovine

serum albumin

Purification: The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions. The

solution is free of unconjugated PE and unconjugated antibody.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to

light. Stable until expiry date (EXP) on label.

Directions for Use: For flow cytometry, the suggested use of this antibody is 5 µL per 1 x 10^6 cells in 100 µL or per 100 µL of


whole blood. It is recommended that the antibody be titrated for optimal performance for each application.

#### Anti-Human CD45 Antibody, Clone HI30, PE

## **Antibodies**



### Data



Flow cytometry analysis of human peripheral blood mononuclear cells (PBMCs) labeled with Anti-Human CD45 Antibody, Clone HI30, PE (filled histogram) or a mouse IgG1, kappa PE isotype control antibody (solid line histogram).

## **Related Products**

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

### References

- 1. Canter RJ et al. (2014) Anti-proliferative but not anti-angiogenic tyrosine kinase inhibitors enrich for cancer stem cells in soft tissue sarcoma. BMC Cancer 14(1): 756. (FC)
- 2. Wang LX et al. (2014) Humanized-BLT mouse model of Kaposi's sarcoma-associated herpesvirus infection. Proc Natl Acad Sci USA 111(8): 3146–51. (FC)
- 3. Despeaux M et al. (2012) Focal adhesion kinase splice variants maintain primitive acute myeloid leukemia cells through altered Wnt signaling. Stem Cells 30(8): 1597–610. (FC)
- 4. Mian, F M et al. (2010) Humanized mice are susceptible to Salmonella typhi infection. Cell Mol Immunol 8(1): 83-7. (FC)
- 5. Jiang Q et al. (2008) FoxP3+CD4+ regulatory T cells play an important role in acute HIV-1 infection in humanized Rag2-/-gammaC-/- mice in vivo. Blood 112(7): 2858–68. (FC)
- 6. Nagano M et al. (2007) Identification of functional endothelial progenitor cells suitable for the treatment of ischemic tissue using human umbilical cord blood. Blood 110(1): 151–60. (FC)
- 7. Ninomiya M et al. (2007) Homing, proliferation and survival sites of human leukemia cells in vivo in immunodeficient mice. Leukemia 21(1): 136–42. (FC, IF, IHC)
- 8. Bourna-ter Steege JCA et al. (2004) Angiogenic profile of breast carcinoma determines leukocyte infiltration. Clin Cancer Res 10(21): 7171–8. (FC, IHC)
- 9. Yamada T et al. (2002) CD45 controls interleukin-4-mediated IgE class switch recombination in human B cells through its function as a Janus kinase phosphatase. J Biol Chem 277(32): 28830–5. (FA/Inhibition)
- 10. Esser MT et al. (2001) Differential incorporation of CD45, CD80 (B7-1), CD86 (B7-2), and major histocompatibility complex class I and II molecules into human immunodeficiency virus type 1 virions and microvesicles: implications for viral pathogenesis and immune regulation. J Virol 75(13): 6173–82. (FC, WR)
- 11. Yoshino N et al. (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (Macaca fascicularis) by using anti-human cross-reactive antibodies. Exp Anim 49(2): 97–110. (FC)
- 12. Kishihara K et al. (1993) Normal B lymphocyte development but impaired T cell maturation in CD45-exon6 protein tyrosine phosphatase-deficient mice. Cell 74(1): 143–56. (FC, IP)
- 13. Knapp W et al. (Eds.). (1989) Leucocyte Typing IV: White cell differentiation antigens. New York: Oxford University Press.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2020 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.