

Anti-Rat CD90 Antibody, Clone OX-7, FITC



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Antibodies

Mouse monoclonal IgG1 antibody against mouse, rat, guinea pig CD90/CD90.1, FITC-conjugated

Catalog #60024FI
#60024FI.1

500 µg 0.5 mg/mL
100 µg 0.5 mg/mL

Product Description

The OX-7 antibody reacts with rat CD90 (Thy-1) and mouse CD90.1 (Thy-1.1), the latter being an allelic form of CD90 expressed by mouse strains AKR/J, PL, and FVB/N. The OX-7 antibody does not react with CD90.2, which is expressed by many mouse strains, including CBA and BALB/c. CD90 is a GPI-linked membrane glycoprotein and member of the immunoglobulin superfamily. The 25 kDa core protein is N-glycosylated at three sites, giving rise to molecules with a range of molecular masses (25 - 37 kDa). In the rat, CD90 is expressed by several cell types, including hematopoietic stem cells, immature B cells, thymocytes and neurons. In mouse strains expressing CD90.1, it is found on early-stage hematopoietic cells in the bone marrow, thymocytes, and circulating mature T cells. The OX-7 antibody has been reported to induce leukocyte activation, glomerular nephritis, apoptosis in glomerular mesangial cells, and vascular permeability.

Target Antigen Name:	CD90/CD90.1
Alternative Names:	Thy-1, Thy-1.1
Gene ID:	21838/24832
Species Reactivity:	Mouse (AKR/J and PL mouse strains), Rat, Guinea Pig, Rabbit
Host Species:	Mouse (BALB/c)
Clonality:	Monoclonal
Clone:	OX-7
Isotype:	IgG1, kappa
Immunogen:	Rat thymocyte CD90 (Thy-1) antigen
Conjugate:	FITC (Fluorescein isothiocyanate)

Applications


Verified:	FC
Reported:	FC
Special Applications:	This antibody clone has been verified for purity assessments of cells isolated from compatible mouse strains with EasySep™ kits, including EasySep™ Mouse T Cell Isolation Kit (Catalog #19851).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Purification:	The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. For product expiry date, contact techsupport@stemcell.com.
Directions for Use:	For flow cytometry, the suggested use of this antibody is $\leq 0.25 \mu\text{g}$ per 1×10^6 cells in 100 μL volume. It is recommended that the antibody be titrated for optimal performance for each application.

Data



(A) Flow cytometry analysis of Sprague-Dawley rat brain cells labeled with Anti-Rat CD90 Antibody, Clone OX-7, FITC (filled histogram) or Mouse IgG1, kappa Isotype Control Antibody, Clone MOPC-21, FITC (Catalog #60070FI) (solid line histogram).

(B) Flow cytometry analysis of Sprague-Dawley rat thymocytes labeled with Anti-Rat CD90 Antibody, Clone OX-7, FITC (filled histogram) or Mouse IgG1, kappa Isotype Control Antibody, Clone MOPC-21, FITC (solid line histogram).

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

- Hiramatsu Y et al. (2010) c-Maf activates the promoter and enhancer of the IL-21 gene, and TGF-beta inhibits c-Maf-induced IL-21 production in CD4+ T cells. *J Leukoc Biol* 87(4): 703–12. (FC)
- Dyer KD et al. (2007) Eosinophils from lineage-ablated Delta dβGATA bone marrow progenitors: the dβGATA enhancer in the promoter of GATA-1 is not essential for differentiation ex vivo. *J Immunol* 179(3): 1693–9. (FC)
- Jeng CJ et al. (1998) Thy-1 is a component common to multiple populations of synaptic vesicles. *J Cell Biol* 140(3): 685–98. (IP, WB)
- Dráberová L et al. (1996) Thy-1-mediated activation of rat mast cells: the role of Thy-1 membrane microdomains. *Immunology* 87(1): 141–8. (FA, WB)
- Ishizu A et al. (1995) Thy-1 induced on rat endothelium regulates vascular permeability at sites of inflammation. *Int Immunol* 7(12): 1939–47. (IP)
- Kawachi H et al. (1992) Epitope-specific induction of mesangial lesions with proteinuria by a MoAb against mesangial cell surface antigen. *Clin Exp Immunol* 88(3): 399–404. (WB)
- Hermans MH & Opstelten D. (1991) In situ visualization of hemopoietic cell subsets and stromal elements in rat and mouse bone marrow by immunostaining of frozen sections. *J Histochem Cytochem* 39(12): 1627–34. (IHC)
- Nakashima I et al. (1991) Evidence of synergy between Thy-1 and CD3/TCR complex in signal delivery to murine thymocytes for cell death. *J Immunol* 147(4): 1153–62. (FA)
- Paul LC et al. (1984) Thy-1.1 in glomeruli of rat kidneys. *Kidney Int* 25(5): 771–7. (Electron Microscopy, IHC)
- Mason DW & Williams AF. (1980) The kinetics of antibody binding to membrane antigens in solution and at the cell surface. *Biochem J* 187(1): 1–20.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2021 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.