

Anti-Human CD33 Antibody, Clone HIM3-4, PE



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Antibodies

Mouse monoclonal IgG1 antibody
against human, chimpanzee CD33,
PE-conjugated

Catalog #60096PE	100 Tests	5 µL/test
#60096PE.1	25 Tests	5 µL/test

Product Description

The HIM3-4 antibody reacts with CD33 (Siglec-3), an ~67 - 75 kDa type 1 transmembrane sialoadhesion protein of the immunoglobulin superfamily. CD33 is expressed on committed myeloid progenitor cells, monocytes, granulocytes, dendritic cells, mast cells, resident macrophages, subsets of B cells, and activated T and NK cells. CD33 is a major marker used to differentiate myeloid from lymphoid or erythroid leukemias. In peripheral blood, it is expressed at high levels on monocytes and at lower levels on granulocytes. CD33 functions as a sialic acid-dependent cell adhesion molecule and mediates protein-glycan and protein-protein interactions between cells. It also has a putative role as an inhibitory receptor during the immune response. HIM3-4 reportedly binds to the extracellular C2 Ig domain of CD33 and does not block subsequent binding of several other anti-CD33 clones, including P67.6, WM53, and WM54. Two isoforms designated CD33M and CD33m have been identified, and among these clones only HIM3-4 recognizes both isoforms.

Target Antigen Name:	CD33
Alternative Names:	gp67, p67, Sialic acid-binding immunoglobulin-like lectin 3, SIGLEC3, Siglec-3
Gene ID:	945
Species Reactivity:	Human, Chimpanzee
Host Species:	Mouse
Clonality:	Monoclonal
Clone:	HIM3-4
Isotype:	IgG1, kappa
Immunogen:	NFMY-9s human cell line
Conjugate:	PE (Phycoerythrin)

Applications


Verified:	FC
Reported:	FACS, FC
Special Applications:	This antibody clone has been verified for purity assessments of cells isolated with EasySep™ Human CD14 Positive Selection Kit II (Catalog #17858).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide, 0.1% gelatin, and 0.2% (w/v) bovine serum albumin
Purification:	The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. For product expiry date, contact techsupport@stemcell.com.
Directions for Use:	For flow cytometry, the suggested use of this antibody is ≤ 5 µL (0.5 µg) per 1 × 10 ⁶ cells in 100 µL. It is recommended that the antibody be titrated for optimal performance for each application.

Data



Flow cytometry analysis of human buffy coat nucleated cells labeled with Anti-Human CD33 Antibody, Clone HIM3-4, PE (filled histogram) or Mouse IgG1, kappa Isotype Control Antibody, Clone MOPC-21, PE (Catalog #60070PE) (solid line histogram).

Related Products

For a complete list of antibodies, including other conjugates, sizes, and clones, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

1. Pérez-Oliva AB et al. (2011) Epitope mapping, expression and post-translational modifications of two isoforms of CD33 (CD33M and CD33m) on lymphoid and myeloid human cells. *Glycobiology* 21(6): 757–70. (FC, ICC, IF)
2. Hernández-Caselles T et al. (2006) A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by alternative splicing. *J Leukoc Biol* 79(1): 46–58. (FC)
3. Robillard N et al. (2005) CD33 is expressed on plasma cells of a significant number of myeloma patients, and may represent a therapeutic target. *Leukemia* 19(11): 2021–2. (FC)
4. Schlossman SF et al. (Eds.). (1995) *Leucocyte Typing V: White cell differentiation antigens*. New York: Oxford University Press.
5. Nakamura Y et al. (1994) Expression of CD33 antigen on normal human activated T lymphocytes. *Blood* 83(5): 1442–3. (FC)
6. Andrews RG et al. (1989) Precursors of colony-forming cells in humans can be distinguished from colony-forming cells by expression of the CD33 and CD34 antigens and light scatter properties. *J Exp Med* 169(5): 1721–31. (FC)

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2022 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.