Antibodies	Anti-Me Clone P	ouse NK1.1 (CD161), K136	STENCELL™ T E C H N O L O G I E S
		onoclonal IgG2a antibody nouse NK1.1 (CD161), nated	Scientists Helping Scientists™   WWW.STEMCELL.COM
Catalog #100-0459	100 µg	0.5 mg/mL	INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

### **Product Description**

The PK136 antibody reacts with murine NK1.1 (CD161), an ~80 kDa homodimeric type 2 transmembrane glycoprotein and C-type lectin receptor expressed on NK cells, NK-T cells, and rare subsets of T cells and monocytes in select strains of mice. NK1.1 is encoded by the KIrb1b and KIrb1c genes, which specify CD161b and CD161c polypeptides, respectively. CD161b is expressed only by Swiss NIH and SJL mice, and CD161c by certain strains such as C57BL, FVB/N, and NZB (but not A, AKR, BALB/c, CBA/J, C3H, C57BR, C58, DBA/1, DBA/2, NOD, SJL, or 129). NK1.1 has functional roles in modulating several processes, including the activation and proliferation of NK cells, induction of interferon- $\gamma$  production, and release of cytotoxic granules. Its expression on the cell surface is specifically upregulated by IL-12. For detection of NK cells in NK1.1- strains, Anti-Mouse CD49b Antibody, Clone DX5 (Catalog #60020) is recommended. DX5 recognizes CD49b (integrin  $\alpha$ 2), another commonly used NK cell marker.

Target Antigen Name:	NK1.1 (CD161)
Alternative Names:	CD161b, CD161c, Klrb1b, Klrb1c, Ly-55, Ly-59, NK-1.1, NKRP1, NKRP1a, NKRP1b
Gene ID:	17059
Species Reactivity:	Mouse (strain-specific)
Host Species:	(C3H x BALB/c) F1 hybrid
Clonality:	Monoclonal
Clone:	PK136
Isotype:	lgG2a, kappa
Immunogen:	NK-1+ cells from mouse spleen and bone marrow
Conjugate:	Unconjugated

## Applications

Verified:	FC
Reported:	CyTOF®, Depletion, FA, FC, ICC, IF, IHC, IP
Special Applications:	This antibody clone has been verified for purity assessments of cells isolated from NK1.1+ mouse strains
	such as C57BL/6, FVB/N, NZB, Swiss NIH, and SJL with EasySep™ kits, including EasySep™ Mouse NK Cell
	Isolation Kit (Catalog #19855).


Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

## Properties

Formulation:	Phosphate-buffered saline, pH 7.2, containing 0.09% sodium azide
Purification:	The antibody was purified by affinity chromatography.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Stable until expiry date (EXP) on label.
Directions for Use:	For flow cytometry, the suggested use of this antibody is $\leq 1 \ \mu g$ per 1 x 10^6 cells in 100 $\mu$ L. It is recommended that the antibody be titrated for optimal performance for each application.



Data



(A) Flow cytometry analysis of C57BL/6 mouse splenocytes (gated on lymphocytes) labeled with Anti-Mouse NK1.1 (CD161) Antibody, Clone PK136, followed by a rat anti-mouse IgG2a antibody, PE and Anti-Mouse CD45 Antibody, Clone 30-F11, APC (Catalog #60030AZ).
(B) Flow cytometry analysis of C57BL/6 mouse splenocytes (gated on lymphocytes) labeled with Mouse IgG2a, kappa Isotype Control Antibody, Clone MOPC-173 (Catalog #60071), followed by a rat anti-mouse IgG2a antibody, PE and Anti-Mouse CD45 Antibody, Clone 30-F11, APC.

# Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

## References

1. Markman JL et al. (2020) Loss of testosterone impairs anti-tumor neutrophil function. Nat Commun 11(1): 1–15. (FC)

2. Kim J-I et al. (2018) CRISPR/Cas9-mediated knockout of Rag-2 causes systemic lymphopenia with hypoplastic lymphoid organs in FVB mice. Lab Anim Res 34(4): 166. (FC)

3. Seshadri S et al. (2017) Bacillus anthracis lethal toxin negatively modulates ILC3 function through perturbation of IL-23-mediated MAPK signaling. PLoS Pathog 13(10): 1–23. (FACS)

4. McAllister CS et al. (2013) TLR3, TRIF, and caspase 8 determine double-stranded RNA-induced epithelial cell death and survival in vivo. J Immunol 190(1): 418–27. (Depletion, IHC)

5. Mitchell AJ et al. (2012) Inflammasome-dependent IFN-γ drives pathogenesis in Streptococcus pneumoniae meningitis. J Immunol 189(10): 4970–80. (FC)

6. Xie X et al. (2010) MHC class I D(k) expression in hematopoietic and nonhematopoietic cells confers natural killer cell resistance to murine cytomegalovirus. Proc Natl Acad Sci USA 107(19): 8754–9. (Depletion, FA/Stimulation, FC)

7. Zandi S et al. (2008) EBF1 is essential for B-lineage priming and establishment of a transcription factor network in common lymphoid progenitors. J Immunol 181(5): 3364–72. (FC)

8. Kulesza J et al. (2006) NK cell depletion and recovery in SCID mice treated with anti-NK1.1 antibody. Folia Histochem Cytobiol 44(2): 93–6. (Depletion) 9. Carnemolla B et al. (2002) Enhancement of the antitumor properties of interleukin-2 by its targeted delivery to the tumor blood vessel extracellular matrix. Blood 99(5): 1659–65. (IHC)

10. Kitaichi N et al. (2002) Diminution of experimental autoimmune uveoretinitis (EAU) in mice depleted of NK cells. J Leukoc Biol 72(6): 1117–21. (Depletion)

11. Kanwar JR et al. (2001) Effects of survivin antagonists on growth of established tumors and B7-1 immunogene therapy. J Natl Cancer Inst 93(20): 1541–52. (Depletion, IHC, IF)

12. Carlyle JR et al. (1999) Mouse NKR-P1B, a novel NK1.1 antigen with inhibitory function. J Immunol 162(10): 5917–23. (FA, FC, IP)

13. Kung SK et al. (1999) The NKR-P1B gene product is an inhibitory receptor on SJL/J NK cells. J Immunol 162(10): 5876–87. (FA, FC, IP)

14. Reichlin A & Yokoyama WM. (1998) Natural killer cell proliferation induced by anti-NK1.1 and IL-2. Immunol Cell Biol 76(2): 143-52. (FA/Stimulation, FC)

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2020 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Canada Inc. CyTOF is a registered trademark of Fluidigm Corporation. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.