

Mouse Pancreatic Organoids

Cryopreserved mouse pancreatic exocrine organoids for establishment of organoid cultures

Catalog #70933

200 Organoids



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Cryopreserved Mouse Pancreatic Organoids provide a convenient way to establish or standardize pancreatic organoid cultures in the laboratory. Each vial contains fragments of mouse pancreatic exocrine organoids sufficient to establish two wells of organoid cultures. The organoids are derived from the pancreas of C57BL/6 mice, cultured in PancreaCult™ Organoid Growth Medium (Mouse; Catalog #06040), and cryopreserved in CryoStor® CS10 (Catalog #07930). Using cryopreserved Mouse Pancreatic Organoids enables establishment of pancreatic organoid cultures without the need to isolate pancreatic ducts from primary tissue, eliminating the need for access to fresh mouse tissue. The organoids can be passaged and expanded using PancreaCult™ Organoid Growth Medium (Mouse) and cryopreserved in CryoStor® CS10. Mouse pancreatic exocrine organoid cultures can be used for research in a variety of fields, including epithelial cell biology, cancer, and metabolism.

Should you intend to use this product for commercial purposes, please contact HUB at www.huborganoids.nl for a commercial use license or for clarification in relation to HUB licensing.

Properties

Storage: Store at -135°C or colder.

Shelf Life: Stable for 5 years from date of manufacture (MFG) on label.

Contains:

- Frozen mouse pancreatic exocrine organoid segments
- CryoStor® CS10

Materials Required but Not Included

PRODUCT NAME	CATALOG #
Bovine serum albumin (BSA)	---
Corning® Matrigel® Growth Factor Reduced (GFR) Basement Membrane Matrix, Phenol Red-Free, LDEV-Free*	Corning 356231*
Costar® 24-Well Flat-Bottom Plate, Tissue Culture-Treated	38017
DMEM/F-12 with 15 mM HEPES	36254
Falcon® Conical Tubes, 15 mL	38009
PancreaCult™ Organoid Growth Medium (Mouse)	06040

*We recommend using Corning® Matrigel® lots containing ≥ 8 mg/mL protein. Lower protein concentrations may affect organoid growth.

Directions for Use

The following instructions are for preparing one cryovial of organoids for plating in 2 wells of a 24-well tissue culture-treated plate. Use sterile technique throughout the protocol.

NOTE: Pre-wet pipette tips with DMEM/F-12 with 15 mM HEPES + 1% BSA (prepared in step 4) before manipulating organoids. This prevents tissue from sticking to the wall of the pipette tip, which significantly decreases organoid yield.

SETUP

1. Place a 24-well tissue culture-treated plate in a 37°C incubator for at least 1 hour. Place a box of sterile 200 µL pipette tips at 2 - 8°C.
2. Thaw ~75 µL of Corning® Matrigel® on ice.

NOTE: Keep Corning® Matrigel® on ice when thawing and handling to prevent it from gelling.

3. Prepare 5 mL of PancreaCult™ Organoid Growth Medium (refer to the Product Information Sheet [PIS] for PancreaCult™). Warm to room temperature (15 - 25°C).

4. For pre-wetting pipette tips, prepare DMEM/F-12 with 15 mM HEPES + 1% BSA (DMEM + BSA) as follows:
 - a. Add 2 mL of sterile 25% BSA to 48 mL of DMEM/F-12 with 15 mM HEPES. Mix thoroughly. Store at room temperature (15 - 25°C) for the duration of the protocol.
 - b. Store the remaining DMEM + BSA at 2 - 8°C for up to 1 month.

5. To a 15 mL conical tube, add 2 mL of DMEM + BSA (prepared in step 4).

NOTE: Transfer cells to this tube immediately after thawing (steps 6 - 8) to avoid a significant reduction in viability.

THAWING ORGANOIDS

6. Place the cryovial of organoids in a 37°C water bath to thaw for 2 - 2.5 minutes. Thawing is complete when the freezing medium becomes liquid. Perform steps 7 - 8 immediately after cells are thawed.

NOTE: Warming the frozen organoids for too long may affect the growth of the organoids in culture. Once thawed, do not re-freeze.

7. Wipe the outside of the cryovial with 70% ethanol or isopropanol before opening.
8. Using a 1 mL pipettor, add 1 mL of DMEM + BSA to the cryovial. Mix the contents by pipetting up and down 4 times. Immediately transfer the contents of the cryovial to the tube prepared in step 5.
9. Wash the inside of the cryovial and inside of the lid with 2 x 1 mL of DMEM + BSA. Add the washes to the organoid suspension from step 8.
10. Centrifuge the organoid suspension at 290 x g for 5 minutes. If there are bubbles on the surface, aspirate these first, then aspirate the remainder of the supernatant.

CULTURING ORGANOIDS IN MATRIGEL® DOMES

11. Remove the 24-well plate from the incubator and 200 µL pipette tips from the fridge and place in a biosafety cabinet.
12. Process the pellet as described below. Work quickly after adding Matrigel® to the pellet to ensure the Matrigel® does not solidify.

NOTE: The 8 wells in the center of a 24-well plate are the most suitable for domes, since their surfaces are the most even. Wells at the edges of the plate are often slightly slanted, resulting in domes touching the wall of the well and flattening out.

 - a. Using a pipette with a cooled, pre-wetted 200 µL tip, add 62 µL of cold Matrigel® to the pellet. Mix by pipetting up and down 5 - 8 times.
 - b. Set the pipette to 30 µL. Add 30 µL of organoid/Matrigel® suspension to each of 2 wells of the warm 24-well plate such that it forms a dome in the middle of the well. Dispense only to the first stop of the pipettor to avoid generating bubbles on top of the dome.
13. Place the lid on the culture plate. Carefully place the plate in an incubator at 37°C and 5% CO₂ for 10 minutes to let domes solidify.
14. Remove the plate from the incubator and place in the biosafety cabinet.
15. Without disturbing the domes, carefully add 750 µL of PancreaCult™ Organoid Growth Medium against the side of each well containing a dome. Do not pipette directly onto the domes.
16. Add sterile PBS to any unused wells. Place the lid on the culture plate.
17. Capture one 2X image per well using a brightfield microscope (Day 0 images). Incubate at 37°C and 5% CO₂.

NOTE: To monitor organoid growth, take photos of the same field of view within each dome every 2 - 3 days until they are passaged.
18. Perform a full-medium change every 2 - 3 days for up to 1 week by carefully aspirating the medium and adding 750 µL of fresh PancreaCult™ Organoid Growth Medium at room temperature (15 - 25°C). Organoids should be ready for passaging after 4 - 7 days.

NOTE: If Matrigel® domes are loose, change medium by removing 250 µL of medium from the well, then add 500 µL of fresh medium.
19. Passage organoids 1 - 2 times before cryopreservation or downstream experiments; for a passaging protocol, refer to the PIS for PancreaCult™, available at www.stemcell.com, or contact us to request a copy. For the first passage, pool the material from both domes plated in Step 12b (as per PancreaCult™ PIS, section B step 8); thereafter, wells can be passaged individually if desired.

NOTE: For additional passaging and cryopreservation protocols, refer to the Technical Bulletin: Mouse Pancreatic Exocrine Organoid Culture: Supplementary Protocols (Document #27088), available at www.stemcell.com, or contact us to request a copy.



This product was developed under a license to intellectual property owned by Hubrecht Organoid Technology (HUB). This product is sold for research use only. Purchase of this product does not include the right to use it for drug screening aiming for commercial gain, equipment validation, biobanking, or for other commercial purposes. Purchasers wishing to use the product for purposes other than basic research use should contact HUB at www.huborganoids.nl to obtain a further license. Purchasers may apply for a License from HUB, which will not be unreasonably withheld by HUB.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED. FOR ADDITIONAL INFORMATION ON QUALITY AT STEMCELL, REFER TO WWW.STEMCELL.COM/COMPLIANCE.

Copyright © 2025 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and PancreaCult are trademarks of STEMCELL Technologies Canada Inc. Corning, Costar, Falcon, and Matrigel are registered trademarks of Corning Incorporated. CryoStor is a registered trademark of BioLife Solutions. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.