

Small Molecules

Y-27632 (Dihydrochloride)

RHO/ROCK pathway inhibitor; Inhibits ROCK1 and ROCK2

Catalog #	72302	1 mg
	72304	5 mg
	72307	5 x 10 mg
	72308	50 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Y-27632 (Dihydrochloride) is a cell-permeable, highly potent and selective inhibitor of Rho-associated, coiled-coil containing protein kinase (ROCK). Y-27632 inhibits both ROCKI ($K_i = 220$ nM) and ROCKII ($K_i = 300$ nM) by competing with ATP for binding to the catalytic site (Davies et al.; Ishizaki et al.).

Alternative Names:	ROCK inhibitor
CAS Number:	129830-38-2
Chemical Formula:	$C_{14}H_{21}N_3O \cdot 2HCl$
Molecular Weight:	320.3 g/mol
Purity:	$\geq 98\%$
Chemical Name:	4-[(1R)-1-aminoethyl]-N-4-pyridinyl-trans-cyclohexanecarboxamide, dihydrochloride
Structure:	



Properties

Physical Appearance:	A crystalline solid
Storage:	Product stable at $-20^{\circ}C$ as supplied. Protect from prolonged exposure to light. For long-term storage, store with a desiccant. Stable as supplied for 12 months from date of receipt.
Solubility:	<ul style="list-style-type: none">· PBS (pH 7.2) ≤ 30 mM· DMSO ≤ 90 mM· Absolute ethanol ≤ 15 mM For example, to prepare a 5 mM stock solution in PBS or water, resuspend 1 mg in 624 μ L of PBS (pH 7.2) or water.

Prepare stock solution fresh before use. Stock solutions in PBS or water are stable at $-20^{\circ}C$ for up to 6 months. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. This product has been shown to be effective at a final concentration of 10 μ M (Ungrin et al.; Watanabe et al.). Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

MAINTENANCE AND SELF-RENEWAL

- Enhances survival of human embryonic stem (ES) cells when they are dissociated to single cells by preventing dissociation-induced apoptosis (anoikis), thus increasing their cloning efficiency (Watanabe et al.).
- Improves embryoid body formation using forced-aggregation protocols (Ungrin et al.).
- Increases the survival of cryopreserved single human ES cells after thawing (Li et al.).
- Blocks apoptosis of mouse ES-derived neural precursors after dissociation and transplantation (Koyanagi et al.).

REPROGRAMMING

- Direct lineage reprogramming of fibroblasts to mature neurons, in combination with CHIR99021 (Catalog #72052), RepSox (Catalog #73792), Forskolin (Catalog #72112), SP600125 (Catalog #72642), Gö6983 (Catalog #72462), and Valproic Acid (Catalog #72292) (Hu et al.).

DIFFERENTIATION

- Improves survival of human ES cell monolayers at the initiation of differentiation protocols (Rezania et al.)

References

- Davies SP et al. (2000) Specificity and mechanism of action of some commonly used protein kinase inhibitors. *Biochem J* 351(Pt 1): 95–105.
- Hu W et al. (2015) Direct conversion of normal and Alzheimer's Disease human fibroblasts into neuronal cells by small molecules. *Cell Stem Cell* 17(2): 204–12.
- Ishizaki T et al. (2000) Pharmacological properties of Y-27632, a specific inhibitor of rho-associated kinases. *Mol Pharmacol* 57(5): 976–83.
- Koyanagi M et al. (2008) Inhibition of the Rho/ROCK pathway reduces apoptosis during transplantation of embryonic stem cell-derived neural precursors. *J Neurosci Res* 86(2): 270–80.
- Li X et al. (2009) ROCK inhibitor improves survival of cryopreserved serum/feeder-free single human embryonic stem cells. *Hum Reprod* 24(3): 580–9.
- Rezania A et al. (2014) Reversal of diabetes with insulin-producing cells derived in vitro from human pluripotent stem cells. *Nat Biotechnol* 32(11): 1121–33.
- Ungrin MD et al. (2008) Reproducible, ultra high-throughput formation of multicellular organization from single cell suspension-derived human embryonic stem cell aggregates. *PLoS One* 3(2): e1565.
- Watanabe K et al. (2007) A ROCK inhibitor permits survival of dissociated human embryonic stem cells. *Nat Biotechnol* 25(6): 681–6.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, visit www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2021 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.