
Small Molecules	1-Oleoyl Lysophosphatidic Acid	STENCELL <sup>M</sup>
	Lysophosphatidic acid (LPA)1 and LPA2 agonist	Scientists Helping Scientists <sup>™</sup>   WWW.STEMCELL.COM
Catalog # 72694	5 mg	TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713
		INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM
		FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

### **Product Description**

1-Oleoyl Lysophosphatidic Acid is a species of lysophosphatidic acid (LPA) containing oleic acid at the *sn*-1 position. LPA binds to four different G-protein-linked receptors (Chun et al.) to mediate a variety of biological responses including cell proliferation, smooth muscle contraction, platelet aggregation, neurite retraction, and cell motility (Moolenaar). 1-Oleoyl Lysophosphatidic Acid is the most potent of the LPA analogs for calcium mobilization in A431 cells (Jalink et al.) and for growth stimulation of a variety of cell lines (van Corven et al.). This product is supplied as the sodium salt of the molecule.

Molecular Name:	1-Oleoyl Lysophosphatidic Acid (Sodium Salt)	
Alternative Names:	Oleoyl-sn-3-glycerophosphate	
CAS Number:	325465-93-8	
Chemical Formula:	$C_{21}H_{40}O_7P\cdot Na$	
Molecular Weight:	458.5 g/mol	
Purity:	≥ 95%	
Chemical Name:	1-O-9Z-Octadecenoyl-sn-glyceryl-3-phosphoric acid sodium salt	
Structure:		
	ОН	



# Properties

Physical Appearance:	A crystalline solid
Storage:	Product stable at -20°C as supplied. Protect from prolonged exposure to light. Stable as supplied for 12 months from date of receipt.
Solubility:	<ul> <li>PBS (pH 7.2) ≤ 18 mM</li> <li>For example, to prepare a 10 mM stock solution in PBS, resuspend 1 mg in 218 µL of PBS (pH 7.2).</li> <li>Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in PBS at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.</li> </ul>
	For use so a call outure supplement stack solution should be diluted into outure medium immediately before

For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use.



## **Published Applications**

DIFFERENTIATION

· Stimulates neuronal differentiation in cultured mouse or rat neural progenitor cells (Cui & Qiao; Fukushima et al.; Spohr et al.).

· Inhibits human embryonic stem (ES) cell-derived neural stem cells (NSCs) from forming neurospheres and differentiating into neurons in vitro (Dottori et al.).

· Stimulates differentiation of human adipose tissue-derived mesenchymal stem cells to myofibroblast-like cells in vitro (Jeon et al.).

#### References

Chun J et al. (2002) International Union of Pharmacology. XXXIV. Lysophospholipid receptor nomenclature. Pharmacol Rev 54(2): 265–9. Cui HL & Qiao JT. (2006) Promotive action of lysophosphatidic acid on proliferation of rat embryonic neural stem cells and their differentiation to cholinergic neurons in vitro. Sheng Li Xue Bao 58(6): 547–55.

Dottori M et al. (2008) Lysophosphatidic acid inhibits neuronal differentiation of neural stem/progenitor cells derived from human embryonic stem cells. Stem Cells 26(5): 1146–54.

Fukushima N et al. (2007) Lysophosphatidic acid stimulates neuronal differentiation of cortical neuroblasts through the LPA1-G(i/o) pathway. Neurochem Int 50(2): 302–7.

Jalink K et al. (1995) Lysophosphatidic acid-induced Ca2+ mobilization in human A431 cells: structure-activity analysis. Biochem J 307 (Pt 2): 609–16.

Jeon ES et al. (2008) Cancer-derived lysophosphatidic acid stimulates differentiation of human mesenchymal stem cells to myofibroblast-like cells. Stem Cells 26(3): 789–97.

Moolenaar WH. (1994) LPA: a novel lipid mediator with diverse biological actions. Trends Cell Biol 4(6): 213-9.

Spohr TCS et al. (2008) Lysophosphatidic acid receptor-dependent secondary effects via astrocytes promote neuronal differentiation. J Biol Chem 283(12): 7470–9.

Van Corven EJ et al. (1992) Mitogenic action of lysophosphatidic acid and phosphatidic acid on fibroblasts. Dependence on acyl-chain length and inhibition by suramin. Biochem J 281 (Pt 1): 163–9.

### **Related Small Molecules**

For a complete list of small molecules available from STEMCELL Technologies, visit www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2021 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.