Cytokines		Human Recombinant EGF	STEMCELL ^M
		Epidermal growth factor	Scientists Helping Scientists™ WWW.STEMCELL.COM
Catalog #	78006.1 78006 78006.2	100 µg 500 µg 1000 µg	TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Epidermal growth factor (EGF) is characterized by high-affinity binding to various EGF receptors (EGFRs) and the production of mitogenic responses (Carpenter & Cohen). EGF promotes EGFR dimerization, resulting in activation of downstream pathways including PI3K, ERK1/2, JAK/STAT, β -catenin, and calcium signaling. EGF is secreted by the gut-associated salivary and Brunner's glands, is found in a variety of body fluids, and stimulates cell proliferation and differentiation in rodent and neonatal human intestine (Wright et al.). Central nervous system stem cells also proliferate in response to the EGF stimulus (Reynolds & Weiss).

Product Information

Beta-urogastrone, HOMG4, Epidermal growth factor, Pro-epidermal growth factor, URG, Urogastrone
P01133
MNSDSECPLS HDGYCLHDGV CMYIEALDKY ACNCVVGYIG ERCQYRDLKW WELR
6.2 kDa
Human
Mouse, Rat, Monkey
Lyophilized after dialysis against phosphate buffer containing sodium chloride, pH 7.0.
E. coli

Specifications

Activity:	The specific activity is $\ge 5 \times 10^{6}$ units/mg (EC50 ≤ 0.2 ng/mL) as determined by a cell proliferation assay of BALB/c 3T3 cells.
Purity:	≥ 95%
Endotoxin Level:	Measured by kinetic Limulus amebocyte lysate (LAL) analysis and is \leq 0.2 EU/µg protein.

Preparation and Storage

Storage:	Store at -80°C.	
Stability:	Stable as supplied for 12 months from date of receipt.	
Preparation:	Centrifuge vial before opening. Reconstitute product in sterile 18 megohm-cm water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at 2 - 8°C for up to 1 month or at -20°C to -80°C for up to 6 months. Avoid repeated freeze-thaw cycles.	

Cytokines



Data



(A) The biological activity of Human Recombinant EGF was tested by its ability to promote the proliferation of BALB/c 3T3 cells.
Cell proliferation was measured using a fluorometric assay method. The EC50 is defined as the effective concentration of the growth factor at which cell proliferation is at 50% of maximum. The EC50 in the above example is 0.1 ng/mL.
(B) 2 µg of Human Recombinant EGF was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant EGF has a predicted molecular mass of 6.2 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

Carpenter G & Cohen S. (1990) Epidermal growth factor. J Biol Chem 265(14): 7709–12.

Reynolds BA & Weiss S. (1996) Clonal and population analyses demonstrate that an EGF-responsive mammalian embryonic CNS precursor is a stem cell. Dev Biol 175(1): 1–13.

Wright NA et al. (1990) Induction of a novel epidermal growth factor-secreting cell lineage by mucosal ulceration in human gastrointestinal stem cells. Nature 343(6253): 82–5.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2019 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.