Cytokines

Human/Mouse Recombinant TGF-beta 3,

Transforming growth factor beta 3, animal component-free

Catalog # 78156 10 μg 0.25 mg/mL

78156.1 100 μg 0.25 mg/mL 78156.2 4 x 250 μg 0.25 mg/mL



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Transforming growth factor (TGF) beta 3 is a member of the TGF-β superfamily and regulates diverse cellular phenotypes. TGF-beta 3 binds to serine-threonine kinase type I and II receptors and activates signal transduction through SMAD2/3 proteins, thus regulating a variety of functions, including cell proliferation, differentiation, wound healing, apoptosis, and metabolism (Massagué; McDowell et al.). TGF-beta 3 enhances glycosaminoglycan production by mesenchymal stromal cells, stimulates scar-free healing, and improves glucose tolerance and phenotypic changes in adipocyte morphology (Hall et al.; Holton et al.). TGF-beta 3 induces proliferation of posterofrontal suture-derived mesenchymal stromal cells, and stimulates expression of fibroblast growth factors 2 and 18 (James et al.). This product is animal component-free.

Product Information

Alternative Names: ARVD, FLJ16571, TGFB3, TGF-b3, Transforming growth factor beta-3

Accession Number: P10600

Amino Acid Sequence: MALDTNYCFR NLEENCCVRP LYIDFRQDLG WKWVHEPKGY YANFCSGPCP YLRSADTTHS TVLGLYNTLN

PEASASPCCV PQDLEPLTIL YYVGRTPKVE QLSNMVVKSC KCS

Predicted Molecular Mass: 12.9 kDa monomer; 25.7 kDa dimer

Species: Human, Mouse

Formulation: Sterile-filtered liquid containing 10 mM acetic acid and 20% ethanol.

Source: E. coli

Specifications

Activity: The specific activity is $\geq 1.0 \times 10^6$ units/mg (EC50 ≤ 1 ng/mL) as determined by the ability to inhibit IL-4-

induced HT-2 cell proliferation.

Purity: $\geq 95\%$

Endotoxin Level: Measured by kinetic Limulus amebocyte lysate (LAL) analysis and is ≤ 1 EU/µg protein.

Preparation and Storage

Storage: Store at 2 - 8°C.

Stability: Stable as supplied for 12 months from date of receipt.

Preparation: Centrifuge vial before opening. Do not vortex. Store at 2 - 8°C for up to 12 months from date of receipt.


Do not freeze.

Human/Mouse Recombinant TGF-beta 3, ACF

Cytokines



Data



- (A) The biological activity of Human/Mouse Recombinant TGF-beta 3, ACF was tested by its ability to inhibit proliferation of HT-2 cells in the presence of human IL-4. The EC50 is defined as the effective concentration of the growth factor at which cell proliferation inhibition is at 50% of maximum. The EC50 in the above example is 0.038 ng/mL.
- (B) 1 µg of Human/Mouse Recombinant TGF-beta 3, ACF was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human/Mouse Recombinant TGF-beta 3, ACF has a predicted molecular mass of 12.9 kDa (monomer) or 25.7 kDa (dimer).

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

Hall BE et al. (2013) Transforming growth factor-β3 (TGF-β3) knock-in ameliorates inflammation due to TGF-β1 deficiency while promoting glucose tolerance. J Biol Chem 288(44): 32074–92.

Holton J et al. (2016) The basic science of bone marrow aspirate concentrate in chondral injuries. Orthop Rev (Pavia) 8(3): 6659. James AW et al. (2009) Differential effects of TGF-beta1 and TGF-beta3 on chondrogenesis in posterofrontal cranial suture-derived mesenchymal cells in vitro. Plast Reconstr Surg 123(1): 31–43.

Massagué J. (2000) How cells read TGF-beta signals. Nat Rev Mol Cell Biol 1(3): 169-78.

McDowell N et al. (1997) Activin has direct long-range signalling activity and can form a concentration gradient by diffusion. Curr Biol 7(9): 671–81.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2021 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.