

Animal Component-Free Cell Dissociation Kit

Dissociation kit for human stem and progenitor cells

Catalog #05426

1 Kit



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Animal Component-Free Cell Dissociation Kit is optimized for dissociation and passaging of human stem and progenitor cells cultured in various media formulations.

Product Information

The following components are sold as a complete kit (Catalog #05426) and are not available for individual sale.

COMPONENT NAME	COMPONENT #	SIZE	STORAGE	SHELF LIFE
ACF Enzymatic Dissociation Solution	05427	250 mL	Store at -20°C.	Stable until expiry date (EXP) on label.
ACF Enzyme Inhibition Solution	05428	250 mL	Store at -20°C.	Stable until expiry date (EXP) on label.

Directions for Use

For dissociation and passaging of human mesenchymal stem and progenitor cells (MSCs), refer to the protocol below. For other cell types, use as directed in the Product Information Sheet (PIS) for the applicable STEMCELL product.

NOTE: For instructions on culturing human MSCs using serum- and animal component-free MesenCult™-ACF Plus Medium (Catalog #05445), refer to the PIS (Document #10000003462), available at www.stemcell.com, or contact us to request a copy.

The following protocol is for passaging MSCs cultured in MesenCult™-ACF Plus Medium in a single T-75 cm² flask (e.g. Catalog #200-0500). If using other cultureware, adjust cell numbers and volumes accordingly.

1. Thaw ACF Enzymatic Dissociation Solution and ACF Enzyme Inhibition Solution at 2 - 8°C. Mix each solution thoroughly.
NOTE: Once thawed, use immediately or store at 2 - 8°C for up to 4 weeks. Alternatively, aliquot and store solutions at -20°C. Do not exceed the shelf life of the solutions. After thawing the aliquoted solutions, use immediately. Do not re-freeze.
2. Warm the ACF Enzymatic Dissociation Solution and ACF Enzyme Inhibition Solution to room temperature (15 - 25°C). Do not incubate at 37°C.
3. Wash cells once with 4 mL of D-PBS (Without Ca⁺⁺ and Mg⁺⁺; Catalog #37350).
4. Add 6 mL of ACF Enzymatic Dissociation Solution.
5. Incubate at 37°C for 2 - 6 minutes. Tap the flask to detach cells. If less than 90% of cells have detached, incubate at 37°C for an additional 1 - 2 minutes and tap the flask again.
6. Add 6 mL of ACF Enzyme Inhibition Solution and collect cells in a polypropylene tube (e.g. 50 mL polypropylene tube [Catalog #200-0520]).
7. Wash the flask with 6 mL of complete MesenCult™-ACF Plus Medium.
8. Place wash into the same polypropylene tube as in step 6.
9. Centrifuge the tube at 300 x g for 8 minutes with the **brake on**.
10. Discard the supernatant and resuspend the cell pellet in desired culture medium.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2023 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and MesenCult are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.