

mTeSR™ Plus

Stabilized feeder-free maintenance medium for human ES and iPS cells

Catalog #05825

1 Kit



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

mTeSR™ Plus is a stabilized, serum-free cell culture medium for the feeder-free maintenance and expansion of human embryonic stem (ES) and induced pluripotent stem (iPS) cells. It is based on the mTeSR™1 formulation^{1,2}, the most widely published feeder-free cell culture medium for human ES and iPS cells.

To enhance cell quality attributes, particularly during restricted feeds, critical medium components have been stabilized, including fibroblast growth factor 2 (FGF2; also known as basic FGF [bFGF]), and medium pH is more consistent. As a result, mTeSR™ Plus allows for greater cell expansion rates with daily feeding, while also maintaining cell quality during restricted feeding schedules.

mTeSR™ Plus is compatible with a variety of culture matrices, including Corning® Matrigel® hESC-Qualified Matrix (Catalog #07181) and Vitronectin XF™ (Catalog #07180, a matrix developed and manufactured by Nucleus Biologics).

Each lot of mTeSR™ Plus 5X Supplement is used to prepare complete mTeSR™ Plus medium and then performance-tested in a culture assay using human pluripotent stem cells (hPSCs).

Product Information

The following components are sold as part of the mTeSR™ Plus kit (Catalog #05825) and are not available for individual sale.

COMPONENT NAME	COMPONENT #	SIZE	STORAGE	SHELF LIFE
mTeSR™ Plus Basal Medium	05826	400 mL	Store at 2 - 8°C.	Stable until expiry date (EXP) on label.
mTeSR™ Plus 5X Supplement	05827	100 mL	Store at -20°C.	Stable until expiry date (EXP) on label.

Preparation of Complete mTeSR™ Plus Medium

Use sterile technique to prepare complete mTeSR™ Plus medium (Basal Medium + 5X Supplement). The following example is for preparing 500 mL of complete medium. If preparing other volumes, adjust accordingly.

NOTE: Thaw supplements or complete medium at room temperature (15 - 25°C) or overnight at 2 - 8°C. **Do not thaw in a 37°C water bath.**

1. Thaw mTeSR™ Plus 5X Supplement at room temperature (15 - 25°C) or overnight at 2 - 8°C. **Warm to room temperature.** Mix thoroughly.

NOTE: mTeSR™ Plus 5X Supplement may appear slightly cloudy after thawing. If this is noted, ensure that the supplement is at room temperature (15 - 25°C). If cloudiness persists, place in a 37°C water bath for approximately 5 minutes, swirling occasionally until supplement becomes clear. Supplement must be free of cloudiness before adding to basal medium (step 2).

NOTE: Once thawed, use supplement immediately or aliquot and store at -20°C for up to 3 months. Do not exceed the shelf life of the supplement. After thawing the aliquoted supplement, use immediately. Do not re-freeze.

2. Add 100 mL of mTeSR™ Plus 5X Supplement to 400 mL of mTeSR™ Plus Basal Medium. Mix thoroughly.

NOTE: If not used immediately, store complete mTeSR™ Plus medium at 2 - 8°C for up to 2 weeks. Alternatively, aliquot and store at -20°C for up to 6 months. Do not exceed the shelf life of the individual components. After thawing the aliquoted complete medium, use immediately or store at 2 - 8°C for up to 2 weeks. Do not re-freeze.

If prepared aseptically, complete mTeSR™ Plus medium is ready for use. If desired, the medium can be filtered using a 0.2 - 0.22 µm low protein binding polyethersulfone (PES) filter unit (e.g. Fisher 09-741-04 [0.2 µm, 250 mL]; Fisher SCGP00525 [0.22 µm, 50 mL]).

Directions for Use

For complete instructions on how to maintain human ES and iPS cells in mTeSR™ Plus, refer to the Technical Manual: Maintenance of Human Pluripotent Stem Cells in mTeSR™ Plus (Document #10000005507), available at www.stemcell.com or contact us to request a copy.

Feeding Schedule

mTeSR™ Plus enables using a more flexible feeding schedule without affecting culture quality. To determine a convenient schedule that suits your lab's routine, follow the table below to determine the volume of medium required at each feed. Any combination of feeding intervals can be used throughout a passage when following these guidelines.

FEEDING INTERVAL		
DAILY FEEDING	SKIP ONE DAY	SKIP TWO CONSECUTIVE DAYS
Standard feed volume (e.g. 2 mL per well of a 6-well plate)	Standard feed volume (e.g. 2 mL per well of a 6-well plate)	Double feed volume (e.g. 4 mL per well of a 6-well plate)

Notes for mTeSR™1 Users

Cultures grown in mTeSR™ Plus are very similar to cultures grown in mTeSR™1. An enhanced growth rate may be observed with mTeSR™ Plus, resulting in larger colonies and higher confluence cultures sooner after passaging. Therefore, experienced mTeSR™1 users may note one or more of the following slight adjustments to the passaging parameters established for mTeSR™1 cultures, as outlined in the Technical Manual for mTeSR™ Plus:

- Increased dissociation time during passaging of larger colonies
- Increased split ratio to maintain similar confluency
- Decreased passaging interval for more rapidly growing cultures

Assessment of hPSCs

The following antibodies can be used to characterize hPSCs by flow cytometry or immunocytochemistry:

- Anti-Human SSEA-4 Antibody, Clone MC-813-70 (Catalog #60062)
- Anti-Human TRA-1-60 Antibody, Clone TRA-1-60R (Catalog #60064)
- Anti-Human OCT4 (OCT3) Antibody, Clone 3A2A20 (Catalog #60093)

For complete flow cytometry protocols and antibodies that can be used, refer to the Technical Manual: Maintenance of Human Pluripotent Stem Cells in mTeSR™ Plus (Document #10000005507), available at www.stemcell.com or contact us to request a copy.

Related Products

For related products, including specialized cell culture and storage media, matrices, antibodies, cytokines, and small molecules, visit www.stemcell.com/hPSCworkflow or contact us at techsupport@stemcell.com.

References

1. Ludwig TE et al. (2006) Derivation of human embryonic stem cells in defined conditions. *Nat Biotechnol* 24(2): 185–7.
2. Ludwig TE et al. (2006) Feeder-independent culture of human embryonic stem cells. *Nat Methods* 3(8): 637–46.



This product was developed under license to intellectual property owned by WiCell™ Research Institute.

This product is sold for research use only (whether the buyer is an academic or for-profit entity) under a non-transferable, limited-use license. Purchase of this product does not include the right to sell, use or otherwise transfer this product for commercial purposes (i.e., any activity undertaken for consideration, such as use of this product for manufacturing, or resale of this product or any materials made using this product, or use of this product or any materials made using this product to provide services) or clinical use (i.e., administration of this product or any material using this product to humans) or the right to implant any material made using this product into an animal by, or in collaboration with, a for-profit entity, for purposes other than basic pre-clinical research applications (including without limitation teratoma assays) to validate the function of the cells. Purchasers wishing to use the product for purposes other than research use should contact Asterias Biotherapeutics, Inc. legal department at (650) 433-2900 or legal@asteriasbio.com. Purchasers who do not agree to the terms and conditions set forth above should return the product in acceptable conditions to the seller for a refund.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2019 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and CellAdhere are trademarks of STEMCELL Technologies Canada Inc. CellAdhere™ Laminin-521 is manufactured by BioLamina. mTeSR is a trademark of WARF. Corning and Matrigel are registered trademarks of Corning Incorporated. Vitronectin XF is developed and manufactured by Nucleus Biologics, and Vitronectin XF is a trademark of Nucleus Biologics. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.