

STEMdiff™ Dorsal and Ventral Forebrain Organoid Differentiation Kits

STEMdiff™ Neural Organoid Maintenance Kit



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog #08620 1 Kit
 Catalog #08630 1 Kit
 Catalog #100-0120 1 Kit

Product Description

STEMdiff™ Dorsal and Ventral Forebrain Organoid Differentiation Kits are matrix-free, serum-free cell culture media that enable the robust generation of human pluripotent stem cell-derived neural organoids. The media work with AggreWell™-generated embryoid bodies to prevent organoid fusion and enable the scalable generation of over 500 highly reproducible organoids per kit. Adapted from protocols by Sergiu Pașca (Birey et al.), these brain-region-specific organoids are three-dimensional in vitro models with a cellular composition and structural organization that is representative of the developing human forebrain. STEMdiff™ Dorsal Forebrain Organoid Differentiation Kit (Catalog #08620) generates tissue of the early developing dorsal pallium, while STEMdiff™ Ventral Forebrain Organoid Differentiation Kit (Catalog #08630) generates tissue of the early developing ventral subpallium. Organoids from different brain regions generated with these kits may be fused to form AssemBloids™ (fused organoids) to allow for advanced neurological modeling. For extended periods of organoid culture (> 50 days), the components required for organoid maintenance are available as STEMdiff™ Neural Organoid Maintenance Kit (Catalog #100-0120).

Ordering Information

PRODUCT NAME	CATALOG #	SIZE	KIT COMPONENTS
STEMdiff™ Dorsal Forebrain Organoid Differentiation Kit	08620	1 Kit	<ul style="list-style-type: none"> • STEMdiff™ Neural Organoid Basal Medium 1 • STEMdiff™ Neural Organoid Basal Medium 2 • STEMdiff™ Neural Organoid Supplement A • STEMdiff™ Neural Organoid Supplement B • STEMdiff™ Neural Organoid Supplement C
STEMdiff™ Ventral Forebrain Organoid Differentiation Kit	08630	1 Kit	<ul style="list-style-type: none"> • STEMdiff™ Neural Organoid Basal Medium 1 • STEMdiff™ Neural Organoid Basal Medium 2 • STEMdiff™ Neural Organoid Supplement A • STEMdiff™ Neural Organoid Supplement B • STEMdiff™ Neural Organoid Supplement C • STEMdiff™ Neural Organoid Supplement D
STEMdiff™ Neural Organoid Maintenance Kit	100-0120	1 Kit	<ul style="list-style-type: none"> • STEMdiff™ Neural Organoid Basal Medium 2 • STEMdiff™ Neural Organoid Supplement A

Component Storage and Stability


The following components are sold as part of a kit (Catalog #08620, 08630, or 100-0120) and are not available for individual sale.

COMPONENT NAME	COMPONENT #	SIZE	STORAGE	SHELF LIFE
STEMdiff™ Neural Organoid Basal Medium 1	08621	20 mL	Store at -20°C.	Stable for 12 months from date of manufacture (MFG) on label.
STEMdiff™ Neural Organoid Basal Medium 2	08622	500 mL	Store at 2 - 8°C.	Stable for 12 months from date of manufacture (MFG) on label.
STEMdiff™ Neural Organoid Supplement A	08623	10 mL	Store at -20°C.	Stable for 12 months from date of manufacture (MFG) on label.
STEMdiff™ Neural Organoid Supplement B	08624	0.25 mL	Store at -20°C.	Stable for 12 months from date of manufacture (MFG) on label.
STEMdiff™ Neural Organoid Supplement C	08625	0.25 mL	Store at -20°C.	Stable for 12 months from date of manufacture (MFG) on label.
STEMdiff™ Neural Organoid Supplement D	08631	0.5 mL	Store at -20°C.	Stable for 12 months from date of manufacture (MFG) on label.

Materials Required But Not Included

PRODUCT NAME	CATALOG #
D-PBS (Without Ca ⁺⁺ and Mg ⁺⁺) (PBS)	37350
DMEM/F-12 with 15 mM HEPES	36254
Gentle Cell Dissociation Reagent	100-0485
Y-27632	72302
Trypan Blue	07050
6-Well Ultra-Low Adherent Plates for Suspension Cultures	27145
AggreWell™800 (1 x 24-well plate)	34811
Anti-Adherence Rinsing Solution	07010
37 µm Reversible Strainer	27215 (Small) OR 27250 (Large)
Wide-bore disposable pipette tips, 200 µL and 1 mL	Fisher Scientific 14-222-730 (200 µL) and 14-222-699 (1 mL)
100 mm Dish, Non-Treated	38045
Conical tubes, 15 mL or 50 mL	e.g. 38009 (15 mL) OR 38010 (50 mL)
Serological pipettes, 5 mL or 10 mL	e.g. 38003 OR 38004

Protocol Diagram



*mTeSR™1, mTeSR™ Plus, or TeSR™-E8™

Preparation of Media

Use sterile technique to prepare STEMdiff™ forebrain organoid media. Prepare each medium as needed in Directions for Use. Refer to Table 1 for medium components, volumes, and in-use storage and stability.

1. Thaw Supplement(s) at room temperature (15 - 25°C). Mix thoroughly.

NOTE: If not used immediately, aliquot Supplement(s) and store at -20°C. Do not exceed the shelf life of the Supplement(s). After thawing aliquots, use immediately. Do not re-freeze.

2. Combine components as indicated in Table 1. Mix thoroughly. Warm to room temperature before use.

NOTE: If not used immediately, store media as indicated in Table 1.

Table 1. Preparation of STEMdiff™ Forebrain Organoid Media (Dorsal Forebrain and Ventral Forebrain)

MEDIUM	COMPONENT	VOLUME	IN-USE STORAGE AND STABILITY
Forebrain Organoid Formation Medium (20 mL)	STEMdiff™ Neural Organoid Basal Medium 1	20 mL	Store at 2 - 8°C for up to 3 weeks. Do not exceed the shelf life of the basal medium.
Seeding Medium (2.5 mL)	STEMdiff™ Neural Organoid Basal Medium 1	2.5 mL	Use immediately.
	Y-27632 (10 µM final concentration)	5 µL of 5 mM stock solution	
Forebrain Organoid Expansion Medium (250 mL)	STEMdiff™ Neural Organoid Basal Medium 2*	245 mL	Store at 2 - 8°C for up to 3 weeks. Do not exceed the shelf life of the basal medium or supplements.
	STEMdiff™ Neural Organoid Supplement A	5 mL	
	STEMdiff™ Neural Organoid Supplement B	0.25 mL	
	STEMdiff™ Neural Organoid Supplement D (For ventral forebrain only)	0.25 mL	
Forebrain Organoid Differentiation Medium (250 mL)	STEMdiff™ Neural Organoid Basal Medium 2*	245 mL	
	STEMdiff™ Neural Organoid Supplement A	5 mL	
	STEMdiff™ Neural Organoid Supplement C	0.25 mL	
Forebrain Organoid Maintenance Medium (100 mL)**	STEMdiff™ Neural Organoid Basal Medium 2*	98 mL	
	STEMdiff™ Neural Organoid Supplement A	2 mL	

*This medium is viscous; pipette slowly to ensure medium is transferred effectively.

**Additional maintenance medium is available for purchase (STEMdiff™ Neural Organoid Maintenance Kit [Catalog #100-0120]).

Directions for Use

Please read the entire protocol before proceeding.

Use sterile technique when performing the following protocols:

- A. Forebrain Organoid Formation (Day 0 - 6)
- B. Forebrain Organoid Expansion (Day 6 - 25)
- C. Forebrain Organoid Differentiation (Day 25 - 43)
- D. Forebrain Organoid Maintenance (Day 43+)

A. FOREBRAIN ORGANOID FORMATION (Day 0 - 6)

The following instructions are for generating a single-cell suspension of human embryonic stem (ES) or induced pluripotent stem (iPS) cells previously cultured in mTeSR™1, mTeSR™ Plus, or TeSR™-E8™ in a 100 mm dish and then plating cells into one well of an AggreWell™800 24-well plate. If using other cultureware, adjust volumes accordingly. Warm cultureware, media, and reagents to room temperature (15 - 25°C) before use.

NOTE: hPSC cultures are ready for passage when the majority of colonies are large, compact, and have dense multi-layered centres. Passage hPSC cultures when they are no more than 70 - 80% confluent and exhibit < 10% differentiation.

Day 0

1. Prepare an AggreWell™800 24-well plate:
 - a. Add 500 µL of Anti-Adherence Rinsing Solution to each well to be used.
 - b. Centrifuge plate at 1300 x g for 5 minutes in a swinging bucket rotor fitted with plate holders.
NOTE: Plates must be well balanced. Prepare a balance plate using a standard plate filled with water to match the weight and position of the AggreWell™ plate.
 - c. Observe plate under a microscope to ensure that bubbles have been removed from microwells.
 - d. If bubbles remain trapped in any microwells, centrifuge at 1300 x g for an additional 5 minutes.
 - e. Aspirate Anti-Adherence Rinsing Solution from the wells.
2. Prepare Seeding Medium (see Preparation of Media) and warm to room temperature.
3. Add 1 mL Seeding Medium to 1 well of the plate prepared in step 1. Set the plate aside.
4. Use a microscope to visually identify regions of differentiation in the hPSC culture. Mark these using a felt tip or lens marker on the bottom of the 100 mm dish. Remove regions of differentiation by scraping with a pipette tip or by aspiration.

- Aspirate medium from hPSC culture and wash the well with 3 - 5 mL sterile phosphate-buffered saline (PBS).
- Aspirate PBS and add 3 mL of Gentle Cell Dissociation Reagent.
- Incubate at 37°C for 8 - 10 minutes.
NOTE: Incubation time may vary when using different cell lines or other non-enzymatic cell dissociation reagents.
- Using a 1 mL pipettor, gently resuspend cells by pipetting up and down slowly 3 - 5 times. Transfer cell suspension to a sterile 15 mL or 50 mL conical tube.
- Rinse the well with an additional 5 - 10 mL of DMEM/F-12 and add this rinse to the tube containing cells.
- Count viable cells using Trypan Blue and a hemocytometer.
- Calculate volume required to obtain 4.5×10^6 total cells (this will be diluted in the next step to obtain a final concentration of 3×10^6 cells/mL).
- Centrifuge cells at $300 \times g$ for 5 minutes.
- Carefully aspirate the supernatant and resuspend cells in 1.5 mL of Seeding Medium (prepared in step 2) to obtain a final concentration of 3×10^6 cells/mL.
- Add 1 mL of single-cell suspension (i.e. 3×10^6 cells) to the well of the AggreWell™800 plate containing Seeding Medium (prepared in step 3). This will result in 10,000 cells/microwell.
NOTE: Ensure that newly plated cells are evenly dispersed across the entire surface of the well by gently pipetting up and down several times.
- Centrifuge the AggreWell™800 plate at $100 \times g$ for 3 minutes. This will capture the cells in the microwells.
NOTE: Plates must be balanced. It is recommended to balance the plate against a standard 24-well plate filled with water to match the weight and position of the AggreWell™800 plate.
- Examine the AggreWell™800 plate under a microscope to ensure that cells are evenly distributed among the microwells.
- Incubate at 37°C.

Day 1 - 5: Partial-Medium Change

NOTE: On Day 1, uniform embryoid bodies (EBs) should be visible in the AggreWell™800 well.

- Carefully remove the AggreWell™800 plate from the incubator, taking care not to disturb the contents.
- Gently remove $2 \times 750 \mu\text{L}$ of the medium from the well using a 1 mL pipettor and discard.
NOTE: Do not disturb the EBs. Keep the pipette tip toward the upper surface of the medium in the well while removing the medium.
- Slowly add $2 \times 750 \mu\text{L}$ of Forebrain Organoid Formation Medium to the well using a 1 mL pipettor.
NOTE: It is important not to disturb the EBs. Do NOT add the medium directly onto the surface of the well. Support the pipette tip by slightly touching the side of the well at the surface level of the remaining medium inside the well. This will allow for a more controlled release of the medium. Release the medium very slowly into the well by setting the Pipette-aid to "gravity" or "slow". Quick release of medium will dislodge the EBs from the wells.
- Incubate at 37°C for 24 hours.
- Repeat steps 18 - 21 on Days 2 - 5.
- On Day 6, proceed to section B for organoid expansion.

B. FOREBRAIN ORGANOID EXPANSION (Day 6 - 25)

NOTE: Warm cultureware, medium, and reagents to room temperature (15 - 25°C) before use.

NOTE: If ultra-low attachment plates are not available, tissue culture-treated cultureware can be used if it is pre-treated with Anti-Adherence Rinsing Solution to prevent cell attachment. Add 1 mL of Anti-Adherence Rinsing Solution to each well, then remove solution from the well and discard.

Day 6

- Prepare Forebrain Organoid Expansion Medium (see Preparation of Media) and warm to room temperature.
- Add 2 mL of Forebrain Organoid Expansion Medium to each well of a 6-Well Ultra-Low Adherent Plate.
NOTE: Aggregates from 1 well of an AggreWell™800 24-well plate can be evenly distributed into the 6-well plate. Do not exceed 50 aggregates per well.
- Place a $37 \mu\text{m}$ Reversible Strainer on top of a 50 mL conical tube. Label the tube "waste".
NOTE: The arrow on the reversible strainer should point upwards. Use a new strainer and a new tube for each AggreWell™800 well to be harvested.
- Remove the medium from the aggregate-containing well and firmly expel it into the well using a 1 mL pipettor with a wide-bore tip. This will dislodge the aggregates from the well.

- Using the same wide-bore tip, aspirate the suspension and filter it through the 37 µm Reversible Strainer. Aggregates will remain on top of the strainer and single cells will flow through into the waste tube.
- Draw up 1 mL of DMEM/F-12 using the wide-bore tip and firmly expel it into the same AggreWell™800 well. While aggregates are in suspension, quickly transfer the suspension into the strainer from step 5.
- Repeat step 6 until all aggregates have been removed from the well. One or two repeats should be sufficient to dislodge all EBs. Examine the well under a microscope to ensure that all aggregates have been removed.
- Invert the strainer over a new 50 mL conical tube and add 2 mL of Forebrain Organoid Expansion Medium onto the strainer to collect all the EBs in the tube.
NOTE: Expansion Medium is viscous, thus a high pipetting force might be required to efficiently collect aggregates from strainer.
- Gently swirl the aggregates to create a suspension and use a wide-bore tip to transfer aggregates into the 6-well plate. Do not exceed 50 aggregates per well of a 6-well plate.
- Gently rock the plate in short, back-and-forth and side-to-side motions to distribute the aggregates across the wells. Visually inspect plate to ensure minimal contact between aggregates.
- Carefully place plate on a level surface in a 37°C incubator. Incubate for 2 days.

Day 8 - 25: Full-medium changes

- Warm Forebrain Organoid Expansion Medium to room temperature.
- Gently tilt the 6-well plate and wait for aggregates to sink to the bottom of the well (~15 - 30 seconds).
- Carefully re-level the plate and use a 1 mL pipettor to remove medium from the top portion of the well.
- Add 2 mL of fresh Forebrain Organoid Expansion Medium to each well. Incubate at 37°C.
- Perform a full-medium change (steps 12 - 15) **every 2 days** on Day 8 - 25.
- On Day 25, proceed to section C for organoid differentiation.

C. FOREBRAIN ORGANOID DIFFERENTIATION (Day 25 - 43)

- Prepare Forebrain Organoid Differentiation Medium (see Preparation of Media) and warm to room temperature (15 - 25°C).
- Gently tilt the 6-well plate and wait for aggregates to sink to the bottom of the well (~15 - 30 seconds).
- Carefully re-level the plate and use a 1 mL pipettor to aspirate medium from the top portion of the well.
- Add 2 mL of fresh Forebrain Organoid Differentiation Medium to each well. Incubate at 37°C.
- Perform a full-medium change (steps 1 - 4) every 2 days on Day 27 - 43.
NOTE: If medium becomes very acidic (bright yellow) on feed days, culture fewer organoids per well (e.g. add < 25 per well of a 6-well plate).
- On Day 43, proceed to section D for organoid maintenance.

D. FOREBRAIN ORGANOID MAINTENANCE (Day 43+)

- Prepare Forebrain Organoid Maintenance Medium (see Preparation of Media) and warm to room temperature (15 - 25°C).
- Gently tilt the 6-well plate and wait for aggregates to sink to the bottom of the well (~15 - 30 seconds).
- Carefully re-level the plate and use a 1 mL pipettor to aspirate medium from the top portion of the well.
- Add 2 mL of fresh Forebrain Organoid Maintenance Medium to each well.
- Perform a full-medium change every 2 - 3 days.
NOTE: If medium becomes very acidic (bright yellow) on feed days, culture fewer organoids per well (e.g. use < 10 per well of a 6-well plate). Once the organoids are larger (i.e. day 50+), increase feed volume to 3 mL per well to ensure they are covered by medium.

Related Products

For related products, including specialized cell culture and storage media, supplements, antibodies, cytokines, and small molecules, visit www.stemcell.com/hPSCNCworkflow or contact us at techsupport@stemcell.com.

References

Birey F et al. (2017) Assembly of functionally integrated human forebrain spheroids. *Nature* 545: 54–59.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED. FOR ADDITIONAL INFORMATION ON QUALITY AT STEMCELL, REFER TO WWW.STEMCELL.COM/COMPLIANCE.

Copyright © 2021 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, AggreWell, and STEMdiff are trademarks of STEMCELL Technologies Canada Inc. mTeSR, TeSR, and E8 are trademarks of WARF. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.