

**Anti-Mouse CD152 (CTLA-4),
Clone UC10-4F10-11, APC**



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Antibodies

Hamster (Armenian) monoclonal IgG1
antibody against mouse CD152 (CTLA-4),
APC-conjugated

Catalog #100-0325
Catalog #100-0326

25 µg 0.2 mg/mL
100 µg 0.2 mg/mL

Product Description

The UC10-4F10-11 antibody reacts with an extracellular epitope on mouse CD152 (CTLA-4), a type I transmembrane glycoprotein receptor expressed on the surface of activated T and B cells and thymocytes. CD152 comprises a disulfide-linked homodimer of ~35 kDa subunits and is a member of the immunoglobulin protein superfamily. It functions as an antagonistic homolog of CD28 by binding the CD28 co-stimulatory ligands, CD80 and CD86. CD152 thereby acts to inhibit CD28-mediated stimulation during the early stages of T cell expansion. It also contributes to the suppressor function of T regulatory cells. CD152 has important roles in immunological tolerance and immunity, and mutations in its cognate gene have been associated with certain autoimmune disorders.

Target Antigen Name:	CD152 (CTLA-4)
Alternative Names:	CTLA-4, Cytotoxic T lymphocyte-associated antigen-4, Ly-56
Gene ID:	12477
Species Reactivity:	Mouse
Host Species:	Hamster (Armenian)
Clonality:	Monoclonal
Clone:	UC10-4F10-11
Isotype:	IgG1, kappa
Immunogen:	Mouse CTLA-4 IgG2a fusion protein
Conjugate:	APC (Allophycocyanin)

Applications


Verified:	FC
Reported:	FC

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation:	Phosphate-buffered saline, pH 7.2, containing 0.09% sodium azide and 0.1% gelatin
Purification:	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. Stable until expiry date (EXP) on label.
Directions for Use:	For flow cytometry, the suggested use of this reagent is $\leq 0.06 \mu\text{g}$ per 1×10^6 cells in 100 μL . It is recommended that the antibody be titrated for optimal performance for each application.

Data



Flow cytometry analysis of 3-day Con A-stimulated C57BL/6 mouse splenocytes labeled with Anti-Mouse CD152 (CTLA-4) Antibody, Clone UC10-4F10-11, APC (filled histogram) or an Armenian hamster IgG, APC isotype control antibody (solid line histogram).

Related Products

For a complete list of antibodies, including other conjugates, sizes, and clones, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

1. Yasuda K et al. (2019) Satb1 regulates the effector program of encephalitogenic tissue Th17 cells in chronic inflammation. *Nat Commun* 10(1): 549. (FC)
2. Rizzo A et al. (2018) ROR γ t-expressing Tregs drive the growth of colitis-associated colorectal cancer by controlling IL6 in dendritic cells. *Cancer Immunol Res* 6(9): 1082–92. (FC)
3. Kishore M et al. (2017) Regulatory T cell migration is dependent on glucokinase-mediated glycolysis. *Immunity* 47(5): 875–89.e10. (FA/Activation)
4. Iraolagoitia XLR et al. (2016) NK cells restrain spontaneous antitumor CD8⁺ T cell priming through PD-1/PD-L1 interactions with dendritic cells. *J Immunol* 197(3): 953–61. (FC)

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2020 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.