Primary Cells

Human Peripheral Blood CD4+CD25+CD127low T Cells, Frozen



TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Scientists Helping Scientists[™] | www.stemcell.com

Product Description

Primary human CD4+CD25+CD127low T cells were isolated from peripheral blood using immunomagnetic separation techniques. CD4+, CD25+, and CD127low are expressed on specific subsets of T cells.

Cells were obtained using Institutional Review Board (IRB)-approved consent forms and protocols.

Donor Status: Normal

Characterization Criteria: Cell count, viability, donor virus testing, age, sex, ethnicity, weight, height

Format: CD4+CD25+CD127low T cells are frozen in CryoStor® CS10. Purity: $\geq 85\%$ CD3+CD4+CD25+CD127low by flow cytometry

Anticoagulant: Acid-citrate-dextrose solution A (ACDA)

For donor details, refer to the lot-specific Certificate of Analysis.

Stability and Storage

Product stable at -135°C or colder for 12 months from date of receipt. Short-term storage of cells (< 1 month) at -80°C is acceptable, but should be minimized to ensure maximum stability. Thawed samples must be used immediately. As these are primary cells, they have a finite life span in culture.

Precautions

Donor Screening: Donors are screened for HIV-1, HIV-2, hepatitis B, and hepatitis C.

If the donor has been screened within 90 days of donation the product will be shipped with negative test results from donor screening.

If the donor has not been screened within 90 days of collection, a test sample will be taken at the time of donation and the product will be shipped before the screening results are available. In the unlikely event that a test result is positive, the customer will be contacted as soon as possible (usually within 24 - 72 hours from the time of shipment).

Donors have been tested and found to be negative for HIV-1, HIV-2, hepatitis B, and hepatitis C prior to donation. As testing cannot completely guarantee that the donor was virus-free, THIS PRODUCT SHOULD BE TREATED AS POTENTIALLY INFECTIOUS and only used following appropriate handling precautions such as those described in biological safety level 2. When handling this product, do not use sharps such as needles and syringes.

STEMCELL cannot guarantee the biological function or any other properties associated with performance of cells in a researcher's individual assay or culture systems. STEMCELL assures the cells will meet the specifications only when assessed, before washing, by our test methods.

FOR IN VITRO RESEARCH USE ONLY. NOT APPROVED FOR DIAGNOSTIC, THERAPEUTIC, OR CLINICAL APPLICATIONS. NOT APPROVED FOR HUMAN OR VETERINARY USE IN VIVO.

Primary Cells

Human Peripheral Blood CD4+CD25+CD127low T Cells, Frozen



Handling / Directions for Use

IMPORTANT: To confirm the number of cells provided, a viable cell count must be done immediately after thawing (before washing). Work quickly once the cells have been thawed to ensure high viability and recovery. Use sterile technique when processing thawed cells.

- 1. Warm medium in a 37°C water bath. See Accessory Products (below) for recommended media.
- 2. Wipe the outside of the vial of cells with 70% ethanol or isopropanol.
- In a biosafety hood, twist the cap a quarter-turn to relieve internal pressure and then retighten.
- Quickly thaw cells in a 37°C water bath by gently shaking the vial. Remove the vial when a small frozen cell pellet remains. Do not vortex
 cells
 - NOTE: It is important to work quickly in the following steps to ensure high cell viability and recovery.
- 5. Wipe the outside of the vial with 70% ethanol or isopropanol.
- 6. Measure the total volume of the cell suspension using a 2 mL serological pipette. This value is used in step 12 to calculate the number of cells provided.
- 7. Remove a 20 µL aliquot of cells for counting. If using Trypan Blue to assess viability, for ≥ 1 x 10^6 cells we suggest adding a minimum of 20 µL of medium and recording the volume of medium added. For < 1 x 10^6 cells, dilute directly in 20 µL Trypan Blue. Set diluted aliquot aside until step 12. See Tips section for more details on performing cell counts with a hemocytometer.</p>
- 8. Transfer the remaining cell suspension to a 50 mL conical tube.
- 9. Rinse the vial with 1 mL of medium and add it dropwise to the cells, while gently swirling the 50 mL tube.
- 10. Wash by adding 15 20 mL of medium dropwise, while gently swirling the tube.
- 11. Centrifuge the cell suspension at 300 x g for 10 minutes at room temperature (15 25°C).
- 12. If using Trypan Blue, perform a cell count on the diluted aliquot from step 7.
- 13. Carefully remove the supernatant (from step 11) with a pipette, leaving a small amount of medium to ensure the cell pellet is not disturbed. Resuspend the cell pellet by gently flicking the tube.
- 14. If cells are starting to clump, add 100 μg DNase I Solution per mL of cell suspension and incubate for 15 minutes at room temperature.
 - NOTE: Do not add DNase I Solution if the cells will be used for DNA or RNA extraction.
- 15. Gently add 15 20 mL of medium to the tube.
- 16. Centrifuge the cell suspension at 300 x g for 10 minutes at room temperature.
- 17. Carefully remove the supernatant with a pipette, leaving a small amount of medium to ensure cell pellet is not disturbed. Resuspend the cell pellet by gently flicking the tube.
 - NOTE: Cell loss of up to 30% can be expected during the wash steps.
- 18. Cells are now ready for use in downstream applications.

Tips

For a protocol on performing total nucleated cell counts using a hemocytometer, refer to https://www.stemcell.com/how-to-count-cells-with-a-hemocytometer.

Accessory Products

PRODUCT NAME	CATALOG #
Falcon® Serological Pipettes, 2 mL	38002
Falcon® Conical Tubes, 50 mL	38010
Iscove's Modified Dulbecco's Medium (add 10% fetal bovine serum)	36150
StemSpan™ SFEM II	09655
Trypan Blue	07050
DNase I Solution (1 mg/mL)	07900

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2022 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Schield Design, StemSpan, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. CryoStor is a registered trademark of BioLife Solutions. Falcon is a registered trademark of Corning Incorporated. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.