

Anti-Human CD154 (CD40L) Antibody, Clone 5C8, APC



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Antibodies

Mouse monoclonal IgG2a antibody against
human CD154 (CD40L), APC-conjugated

Catalog #100-1350

100 Tests 5 µL/test

Product Description

This mouse monoclonal antibody (clone 5C8, APC-conjugated) reacts with human CD154 (CD40L), a member of the tumor necrosis factor alpha (TNF- α) superfamily with a molecular weight of 39 kDa. CD154, also known as CD40L, is a type II transmembrane glycoprotein that is predominantly expressed on activated T-cells but is also found in NK cells, mast cells, basophils, and eosinophils. CD154 (CD40L) binds to CD40, a costimulatory protein found on antigen-presenting cells, such as B cells, monocytes, dendritic cells, and fibroblasts. The CD154 (CD40L)-CD40 interaction induces the activation and proliferation of B-cells, allowing for antibody production, isotype switching, and B-cell memory development. CD154 (CD40L) expression is upregulated in inflammatory bowel disease, suggesting its role in regulating pathogenic cytokine production in T-cells. In vitro studies also demonstrate that the binding of CD40 to its ligand induces endothelial cell activation, highlighting its immunopathogenic role. The CD154 (CD40L), clone 5C8 antibody has been reported to block the interaction and ligation of CD40 with its ligand.

Target Antigen Name:	CD154
Alternative Names:	CD40L, gp39, Ly-62, T-BAM, TRAP
Gene ID:	959
Species Reactivity:	Human
Host Species:	Mouse
Clonality:	Monoclonal
Clone:	5C8
Isotype:	IgG2a, kappa
Immunogen:	Human D1.1 T cell line
Conjugate:	APC (Allophycocyanin)

Applications

Verified: FC

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IHC-P: Immunohistochemistry (paraffin-embedded); IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation:	Phosphate-buffered saline, pH 7.2, containing 0.09% sodium azide and 0.1% gelatin
Purification:	The antibody was purified by affinity chromatography.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. Stable until expiry date (EXP) on label.
Directions for Use:	For flow cytometry, the suggested use of this antibody is 5 µL per 1 x 10 ⁶ cells in 100 µL. It is recommended that the antibody be titrated for optimal performance for each application.

Related Products

For a complete list of antibodies, including other conjugates, sizes, and clones, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/antibodies, or contact us at techsupport@stemcell.com.

References

1. Carbone E, et al. (1997). A new mechanism of NK cell cytotoxicity activation: the CD40–CD40 ligand interaction. *The Journal of Experimental Medicine* 185(12): 2053–60.
2. Grdisa, M. (2003). Influence of CD40 ligation on survival and apoptosis of B-CLL cells in vitro. *Leukemia Research* 27(10): 951–6.
3. Lederman S, et al. (1992) Identification of a novel surface protein on activated CD4+ T cells that induces contact-dependent B cell differentiation (help). *The Journal of Experimental Medicine* 175(4): 1091–101.
4. Liu Z, et al. (1999). Hyperexpression of CD40 ligand (CD154) in inflammatory bowel disease and its contribution to pathogenic cytokine production. *The Journal of Immunology* 163(7): 4049–57.
5. Schuler W, et al. (2004). Efficacy and safety of ABI793, a novel human anti-human CD154 monoclonal antibody, in cynomolgus monkey renal allotransplantation. *Transplantation* 77(5): 717–26.
6. Yellin M. J, et al. (1995). Functional interactions of T cells with endothelial cells: the role of CD40L-CD40-mediated signals. *The Journal of Experimental Medicine* 182(6): 1857–64.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2023 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.