

Anti-Mouse CD4 Antibody, Clone GK1.5

Rat monoclonal antibody against mouse, hamster CD4, unconjugated

Catalog #100-1601

100 µg

0.5 mg/mL

Product Description

This monoclonal antibody (clone GK1.5) reacts with mouse cluster of differentiation 4 (CD4), a 55 kDa single chain type 1 transmembrane glycoprotein, belonging to the immunoglobulin (Ig) superfamily. CD4 contains four extracellular Ig-like domains; D1 - D4. CD4 is expressed at relatively high levels by most thymocytes and a subpopulation of T cells; T helper/inducer cells and at lower levels on dendritic cells. Unlike humans, CD4 is not expressed by murine monocytes or macrophages. CD4 increases the affinity in the interaction between T cell receptor (TCR) and major histocompatibility complex II (MHC II) antigen complex by binding to a non-polymorphic region of MHC II and acting as a co-receptor to TCR in MHC II-restricted antigen recognition. CD4 also amplifies signals from TCR to the cytoplasm through the interaction of its intracellular domain with the cytoplasmic tyrosine kinases such as lymphocyte-specific protein tyrosine kinase (Lck). This monoclonal antibody is widely used as a phenotypic marker for CD4 expression. The RM4-5 clone can block binding of the GK1.5 antibody clone to CD4 on the cell surface. The GK1.5 antibody clone is also reported to be cross-reactive with Syrian hamster CD4.

Target Antigen:	CD4
Alternative Names:	L3T4, T4
Gene ID:	12504
Species Reactivity:	Mouse, Hamster
Host Species:	Rat
Clonality:	Monoclonal
Clone:	GK1.5
Isotype:	IgG2b, kappa
Immunogen:	Mouse CTL, clone V4
Conjugate:	Unconjugated

Applications

Verified Applications: FC

Reported Applications: FC, IHC/IF

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; FCXM: Flow cytometric crossmatch assay; FISH: Fluorescence in situ hybridization; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IHC-F: Immunohistochemistry (frozen-tissue); IHC-P: Immunohistochemistry (paraffin-embedded); IP: Immunoprecipitation; NMR: Nuclear magnetic resonance spectroscopy; RIA: Radioimmunoassay; WB: Western blotting

Properties

Product Formulation: Phosphate-buffered saline, pH 7.2, containing 0.09% sodium azide

Purification: The antibody was purified by affinity chromatography.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. Stable until expiry date (EXP) on label.

Directions for Use: For flow cytometry, the suggested use of this antibody is $\leq 1 \mu\text{g}$ per 1×10^6 cells in 100 μL . It is recommended that the antibody be titrated for optimal performance for each application.

Related Products

For a complete list of antibodies, including other conjugates, sizes, and clones, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/antibodies, or contact us at techsupport@stemcell.com.

References

- Arora S et al. (2006) Effect of a CD4-depleting antibody on the development of cryptococcus neoformans-induced allergic bronchopulmonary mycosis in mice. *Infect Immun* 74(7): 4339.
- Freise AC et al. (2017) ImmunoPET imaging of murine CD4+ T cells using anti-CD4 cys-diabody: Effects of protein dose on T cell function and imaging. *Mol Imaging Biol* 19(4): 599–609.
- Furuta Y et al. (2017) E-NPP3 controls plasmacytoid dendritic cell numbers in the small intestine. *PLoS One* 12(2): e0172509.
- Garo LP et al. (2021) MicroRNA-146a limits tumorigenic inflammation in colorectal cancer. *Nat Commun* 12(1): 2419.
- König R et al. (1992) MHC class II interaction with CD4 mediated by a region analogous to the MHC class I binding site for CD8. *Nature* 356 (6372): 796–8.
- Liu J & Yin J. (2021) Immunotherapy with recombinant Alt a 1 suppresses allergic asthma and influences T follicular cells and regulatory B cells in mice. *Front Immunol* 12: 747730.
- Mayer CT et al. (2013) CD4 blockade directly inhibits mouse and human CD4+ T cell functions independent of Foxp3+ Tregs. *J Autoimmun* 47: 73–82.
- Neumann AK et al. (2005) Hypoxia inducible factor 1 α regulates T cell receptor signal transduction. *Proc Natl Acad Sci U S A* 102(47): 17071.
- Plaks V et al. (2015) Adaptive immune regulation of mammary postnatal organogenesis. *Dev Cell* 34(5): 493–504.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2024 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.