

Anti-Mouse TCR Beta Antibody, Clone H57-597, APC

Hamster (Armenian) monoclonal antibody against mouse T cell receptor beta, APC-conjugated

Catalog #100-1611

100 µg

0.2 mg/mL

Product Description

This monoclonal antibody reacts with the beta chain of mouse T cell receptor (TCR). TCR beta is a member of the immunoglobulin superfamily and together with the TCR alpha chain form the alpha-beta TCR. TCR beta is expressed by thymocytes in a developmentally regulated manner and by majority of peripheral T cells. Small number of T cells express an alternative heteromer of gamma and delta protein chains, known as the gamma-delta TCR. TCR-CD3 complex in conjunction with co-receptors CD4 or CD8 enable the recognition and response to antigens attached to major histocompatibility complex (MHC) molecules on antigen-presenting cells. Crosslinking of the TCR complex with H57-597 induces activation and proliferation of T cells or apoptosis, based on assay conditions. The H57-597 antibody clone has been widely used as a phenotypic indicator for T cells expressing alpha-beta TCR. This antibody does not cross-react with cells expressing the gamma-delta TCR.

Target Antigen:	TCR beta
Alternative Names:	Tcrb, TCR-b chain, TCR beta, Tib
Gene ID:	21577
Species Reactivity:	Mouse
Host Species:	Hamster
Clonality:	Monoclonal
Clone:	H57-597
Isotype:	Armenian hamster IgG
Immunogen:	Affinity-purified TCR from mouse T cell hybridoma DO-11.10
Conjugate:	APC (Allophycocyanin)

Applications

Verified Applications: FC

Reported Applications: FC

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; FCXM: Flow cytometric crossmatch assay; FISH: Fluorescence in situ hybridization; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IHC-F: Immunohistochemistry (frozen-tissue); IHC-P: Immunohistochemistry (paraffin-embedded); IP: Immunoprecipitation; NMR: Nuclear magnetic resonance spectroscopy; RIA: Radioimmunoassay; WB: Western blotting

Properties

Product Formulation: Phosphate-buffered saline, pH 7.2, containing 0.09% sodium azide and 0.1% gelatin

Purification: The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. Stable until expiry date (EXP) on label.

Directions for Use: For flow cytometry, the suggested use of this antibody is $\leq 1 \mu\text{g}$ per 1×10^6 cells in 100 μL . It is recommended that the antibody be titrated for optimal performance for each application.

Related Products

For a complete list of antibodies, including other conjugates, sizes, and clones, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/antibodies, or contact us at techsupport@stemcell.com.

References

- Berent-Maoz B et al. (2012) Fibroblast growth factor-7 partially reverses murine thymocyte progenitor aging by repression of Ink4a. *Blood* 119 (24): 5715–21.
- Doisne J-M et al. (2009) Skin and peripheral lymph node invariant NKT cells are mainly retinoic acid receptor-related orphan receptor γt^+ and respond preferentially under inflammatory conditions. *J Immunol* 183(3): 2142–9.
- Harada N et al. (2004) IL-12 gene therapy is an effective therapeutic strategy for hepatocellular carcinoma in immunosuppressed mice. *J Immunol* 173(11): 6635–44.
- Kaczmarek J et al. (2014) Liver sinusoidal endothelial cell-mediated CD8 T cell priming depends on co-inhibitory signal integration over time. *PLoS One* 9(6): e99574.
- Kisielow J et al. (2019) Deciphering CD4+ T cell specificity using novel MHC–TCR chimeric receptors. *Nat Immunol* 20(5): 652–62.
- Kubo RT et al. (1989) Characterization of a monoclonal antibody which detects all murine alpha beta T cell receptors. *J Immunol* 142(8): 2736–42.
- Matei IR et al. (2007) ATM deficiency disrupts Tcr α locus integrity and the maturation of CD4+CD8+ thymocytes. *Blood* 109(5): 1887–96.
- O'Brien RL et al. (2009) Protective role of gammadelta T cells in spontaneous ocular inflammation. *Invest Ophthalmol Vis Sci* 50(7): 3266–74.
- Oh-hora M et al. (2019) Stromal interaction molecule deficiency in T cells promotes spontaneous follicular helper T cell development and causes type 2 immune disorders. *J Immunol* 202(9): 2616–27.
- Pastor S et al. (2006) Analyses of TCR clustering at the T cell–antigen-presenting cell interface and its impact on the activation of naive CD4+ T cells. *Int Immunol* 18(11): 1615–25.
- Sun D et al. (2001) Myelin antigen-specific CD8+ T cells are encephalitogenic and produce severe disease in C57BL/6 mice. *J Immunol* 166(12): 7579–87.
- Wang D et al. (2012) Inhibition of S-phase kinase-associated protein 2 (Skp2) reprograms and converts diabetogenic T cells to Foxp3 + regulatory T cells. *Proc Natl Acad Sci U S A* 109(24): 9493–8.
- Yumimoto K et al. (2015) F-box protein FBXW7 inhibits cancer metastasis in a non-cell-autonomous manner. *J Clin Invest* 125(2): 621–35.

Anti-Mouse TCR Beta Antibody, Clone H57-597, APC

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2024 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.