Primary Cells

Human Cord Blood CD19+ B Cells, Frozen

Catalog # 70013 1 x 10^6 cells

70013.1 2.5 x 10⁶ cells 70013.2 5 x 10⁶ cells



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Primary human CD19+ B cells were isolated from umbilical cord blood (CB) mononuclear cells (MNCs) using positive immunomagnetic separation techniques. CD19 is primarily expressed on B cells.

Cells were obtained using Institutional Review Board (IRB) approved consent forms and protocols.

Donor Status: Normal

Characterization Criteria: Cell count, viability, purity, donor virus testing, age, sex, ethnicity, weight, height, smoking status, other

information

Format: CD19+ B cells are frozen in CryoStor® CS10.

Purity: The purity of CD19+ B cells is \geq 90% by flow cytometry.

Anticoagulant: Citrate-phosphate-dextrose (CPD)

For donor details, refer to the lot-specific Certificate of Analysis.

Stability and Storage

Product stable at -135°C or colder for 12 months from date of receipt. Short-term storage of cells (< 1 month) at -80°C is acceptable, but should be minimized to ensure maximum stability. Thawed samples must be used immediately. As these are primary cells, they have a finite life span in culture.

Precautions

Donor Screening: Testing for HIV-1 and 2, Hepatitis B, and Hepatitis C is performed on a maternal blood sample and/or on a sample of the donated cord blood.

Cryopreserved products are shipped with negative test results from donor screening.

Donors have been tested and found to be negative for HIV-1 and 2, Hepatitis B, and Hepatitis C prior to donation. As testing cannot completely guarantee that the donor was virus-free, THIS PRODUCT SHOULD BE TREATED AS POTENTIALLY INFECTIOUS and only used following appropriate handling precautions such as those described in biological safety level 2.

Storage of frozen cell products in the vapor phase of a liquid nitrogen storage tank is recommended. Storage in the liquid phase can result in cross contamination if the vial breaks or is not sealed properly. Storage in the liquid phase also increases the potential for liquid nitrogen to penetrate the vial and cause it to explode when removed from storage. Use of a face shield is required as a safety precaution when transferring cells from one container to another. When handling this product do not use sharps such as needles and syringes.

STEMCELL cannot guarantee the biological function or any other properties associated with performance of cells in a researcher's individual assay or culture systems. STEMCELL assures the cells will meet the specifications only when assessed immediately after thawing (before washing) by our test methods.

FOR IN VITRO RESEARCH USE ONLY. NOT APPROVED FOR DIAGNOSTIC, THERAPEUTIC, OR CLINICAL APPLICATIONS. NOT APPROVED FOR HUMAN OR VETERINARY USE IN VIVO.

Primary Cells

Human Cord Blood CD19+ B Cells, Frozen



Handling / Directions for Use

IMPORTANT: To confirm the number of cells provided, a viable cell count must be done immediately after thawing (before washing). Work quickly once the cells have been thawed to ensure high viability and recovery. Use sterile techniques when processing thawed cells.

- 1. Warm medium in a 37°C water bath. See Accessory Products (below) for recommended wash media.
- 2. Wipe the outside of the vial of cells with 70% ethanol or isopropanol.
- 3. In a biosafety hood, twist the cap a quarter-turn to relieve internal pressure and then retighten.
- 4. Quickly thaw cells in a 37°C water bath by gently shaking the vial. Remove the vial when a small frozen cell pellet remains. Do not vortex cells.
 - NOTE: It is important to work quickly in the following steps to ensure high cell viability and recovery.
- 5. Wipe the outside of the vial with 70% ethanol or isopropanol.
- 6. Measure the total volume of the cell suspension using a 2 mL serological pipette. This value is used in step 12 to calculate the number of cells provided.
- 7. Remove a 20 μL aliquot of cells for counting. If using Trypan Blue to assess viability, for ≥ 1 x 10⁶ cells we suggest adding a minimum of 20 μL of medium and recording the volume of medium added. For < 1 x 10⁶ cells dilute cells directly in 20 μL Trypan Blue. Set diluted aliquot aside until step 12. See Tips section for more details on performing cell counts with a hemocytometer.
- 8. Transfer the remaining cell suspension to a 50 mL conical tube.
- 9. Rinse the vial with 1 mL of medium and add it drop wise to the cells, while gently swirling the 50 mL tube.
- 10. Wash by adding 15 20 mL of medium drop wise, while gently swirling the tube.
- 11. Centrifuge the cell suspension at 300 x g for 10 minutes at room temperature (15 25°C).
- 12. If using Trypan Blue, dilute the aliquot from step 7 (if not previously done) during centrifugation, and count.
- 13. Carefully remove the supernatant with a pipette, leaving a small amount of medium to ensure the cell pellet is not disturbed. Resuspend the cell pellet by gently flicking the tube.
- 14. If cells are starting to clump add 100 μg of DNase I Solution per mL of cell suspension and incubate for 15 minutes at room temperature (15 25°C).
 - NOTE: Do not add DNase I Solution if the cells will be used for DNA or RNA extraction.
- 15. Gently add 15 20 mL of medium to the tube.
- 16. Centrifuge the cell suspension at 300 x g for 10 minutes at room temperature (15 25°C).
- 17. Carefully remove the supernatant with a pipette, leaving a small amount of medium to ensure cell pellet is not disturbed. Resuspend the cell pellet by gently flicking the tube.
 - NOTE: Cell loss of up to 30% can be expected during the wash steps.
- 18. Cells are now ready for use in downstream applications.

Tips

Manual cell counting protocol: http://www.humanimmunologyportal.com/protocols/performing-cell-counts-with-a-hemacytometer/ Converting g to rpm: http://www.humanimmunologyportal.com/hiptools/apps/

Accessory Products

PRODUCT NAME	CATALOG #
Anti-Human CD19 Antibody, Clone HIB19 (add 10% fetal bovine serum)	60005

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.