# Anti-Human CD8a Antibody, Clone RPA-T8

### **Antibodies**

Mouse monoclonal IgG1 antibody against human, rhesus, cynomolgus

CD8a, unconjugated

Catalog #60022 100 µg 0.5 mg/mL



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

### **Product Description**

The RPA-T8 antibody reacts with CD8a, a 32 - 34 kDa type I transmembrane glycoprotein which is a subunit of CD8. CD8 is a disulfide-bonded dimer, found either as a heterodimer of CD8a ( $\alpha$ ) and CD8b ( $\beta$ ) subunits (i.e.,  $\alpha\beta$ ) or a homodimer ( $\alpha\alpha$ ). CD8 acts as a co-receptor to the T cell receptor (TCR) during T cell activation by binding MHC Class I molecules presented by an antigen-presenting cell. It functions to strengthen the association between the TCR and MHC I-antigen complex, and to amplify signals from the TCR to the cytoplasm through the interaction of its intracellular domain with cytoplasmic tyrosine kinases such as Lck. The CD8a chain binds to the  $\alpha$ 3 domain of class I MHC molecules. CD8 is expressed by a majority of thymocytes, a subset of mature peripheral blood T cells (T cytotoxic cells), a proportion of  $\beta$  T cells, and at lower levels by NK cells (which predominantly express CD8a homodimers).

Target Antigen Name: CD8a
Alternative Names: Leu2, T8
Gene ID: 925

Species Reactivity: Human, Rhesus, Cynomolgus, Baboon, Chimpanzee, Pigtailed Macaque, Sooty Mangabey

Host Species: Mouse
Clonality: Monoclonal
Clone: RPA-T8
Isotype: IgG1, kappa

Immunogen: Full-length human CD8 protein

Conjugate: Unconjugated

# **Applications**

Verified: CellSep, FC

Reported: CyTOF®, FA, FC, IHC

Special Applications: This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including

EasySep<sup>™</sup> Direct Human CD8+ T Cell Isolation Kit (Catalog #19663), EasySep<sup>™</sup> Human CD8+ T Cell Enrichment Kit (Catalog #19053), EasySep<sup>™</sup> Human CD8 Positive Selection Kit (Catalog #18053), EasySep<sup>™</sup>

Human CD3 Positive Selection Kit (Catalog #18051), EasySep™ HLA CD2 Positive Selection Kit

(Catalog #18657HLA), and EasySep™ HLA T Cell Enrichment Kit (Catalog #19051HLA).

(Satisfy "1866/1127), and Easysop Tiezy Son Embranding "1866/1127).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

# **Properties**

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

Purification: The antibody was purified by affinity chromatography.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. For product expiry date, please contact

techsupport@stemcell.com.

Directions for Use: For flow cytometry the suggested use of this antibody is ≤ 2 µg per 1 x 10^6 cells in 100 µL volume. It is


recommended that the antibody be titrated for optimal performance for each application.

#### Anti-Human CD8a Antibody, Clone RPA-T8

# **Antibodies**



### Data



Flow cytometry analysis of human peripheral blood mononuclear cells (PBMCs) labeled with Anti-Human CD8a Antibody, Clone RPA-T8, followed by Goat Anti-Mouse IgG (H+L) Antibody, Polyclonal, FITC (Catalog #60138FI) (filled histogram), or a mouse IgG1, kappa isotype control antibody, followed by Goat Anti-Mouse IgG (H+L) Antibody, Polyclonal, FITC (solid line histogram).

### **Related Products**

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

### References

- 1. Rout N et al. (2010) Paucity of CD4+ natural killer T (NKT) lymphocytes in sooty mangabeys is associated with lack of NKT cell depletion after SIV infection. PLoS One 5(3): e9787. (FC)
- 2. Kmieciak M et al. (2009) Human T cells express CD25 and Foxp3 upon activation and exhibit effector/memory phenotypes without any regulatory/suppressor function. J Transl Med 7: 89. (FC)
- 3. Thakral D et al. (2008) Differential expression of the human CD8beta splice variants and regulation of the M-2 isoform by ubiquitination. J Immunol 180(11): 7431–42. (FC)
- 4. Mack CL et al. (2004) Biliary atresia is associated with CD4+ Th1 cell-mediated portal tract inflammation. Pediatr Res 56(1): 79-87. (IHC)
- 5. Yoshino N et al. (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (Macaca fascicularis) by using anti-human cross-reactive antibodies. Exp Anim 49(2): 97–110. (FC)
- 6. Kishimoto T et al. (Eds.). (1998) Leucocyte Typing VI: White cell differentiation antigens (pp. 1114-5). New York: Garland Publishing Inc.
- 7. Barclay AN et al. (Eds.). (1997) The Leucocyte Antigen Factsbook, Second Edition (pp. 149-51). New York: Academic Press.
- 8. Sopper S et al. (1997) Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. Cytometry 29(4): 351–62.
- 9. Schlossman S et al. (Eds.). (1995) Leucocyte Typing V: White cell differentiation antigens. New York: Oxford University Press.
- 10. Reimann KA et al. (1994) Use of human leukocyte-specific monoclonal antibodies for clinically immunophenotyping lymphocytes of rhesus monkeys. Cytometry 17(1): 102–8.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists and EasySep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.