

Anti-Human CD51 Antibody, Clone NKI-M9

Antibodies

Mouse monoclonal IgG2a antibody
against human CD51 (integrin α V),
unconjugated



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog #60043

100 μ g 0.5 mg/mL

Product Description

The NKI-M9 antibody reacts with CD51 (vitronectin receptor α chain, or integrin α V), a type I transmembrane glycoprotein containing two subunits (125 kDa and 24 kDa) which are generated by post-translational cleavage and are linked by a disulfide bond. CD51 associates non-covalently with integrin β 1 (CD29), β 3 (CD61), β 5, β 6, or β 8 to form heterodimeric cell adhesion receptors for extracellular matrix components such as fibrinogen, collagen, fibronectin, laminin, osteopontin, thrombospondin, vitronectin, and von Willebrand factor. For example, association of CD51 and CD61 forms the integrin α V/ β 3 receptor primarily involved in binding vitronectin. In addition to mediating adhesion and cytoskeletal organization, CD51-containing integrins have roles in signal transduction and thereby modulate processes such as cell proliferation, differentiation, and migration. Both ligand binding and ligand-induced receptor clustering are necessary for initiating integrin-mediated responses. CD51 is expressed broadly on many types of cells, including endothelial cells, fibroblasts, monocytes, macrophages, platelets (at relatively low levels), and osteoclasts. It is also found on hepatoma, melanoma, and neuroblastoma cells.

Target Antigen Name:	CD51 (Integrin α V)
Alternative Names:	α V integrin, integrin α V, vitronectin receptor α chain
Gene ID:	3685
Species Reactivity:	Human
Host Species:	Mouse
Clonality:	Monoclonal
Clone:	NKI-M9
Isotype:	IgG2a, kappa
Immunogen:	Human melanoma cells
Conjugate:	Unconjugated

Applications


Verified:	FC
Reported:	ELISA, FA, FC, IP
Special Applications:	This antibody clone has been verified for labeling human mesenchymal cells grown in MesenCult™ Proliferation Kit (Human; Catalog #05411) and MesenCult™-XF Medium (Catalog #05420).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Purification:	The antibody was purified by affinity chromatography.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. For product expiry date, please contact techsupport@stemcell.com .
Directions for Use:	For flow cytometry the suggested use of this antibody is $\leq 2 \mu$ g per 1×10^6 cells in 100 μ L volume. It is recommended that the antibody be titrated for optimal performance for each application. Clone NKI-M9 is reportedly not suitable for IHC with paraffin-embedded sections.

Data



Flow cytometry analysis of HT1080 human fibrosarcoma cells labeled with Anti-Human CD51 Antibody, Clone NKI-M9, followed by Goat Anti-Mouse IgG (H+L) Antibody, Polyclonal, FITC (Catalog #60138FI) (filled histogram), or Mouse IgG2a, kappa Isotype Control Antibody, Clone MOPC-173 (Catalog #60071), followed by Goat Anti-Mouse IgG (H+L) Antibody, Polyclonal, FITC (solid line histogram).

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

1. Zola H et al. (2007) *Leukocyte and Stromal Cell Molecules: The CD Markers*. New York: John Wiley & Sons Inc.
2. Burdick MM et al. (2003) Colon carcinoma cell glycolipids, integrins, and other glycoproteins mediate adhesion to HUVECs under flow. *Am J Physiol Cell Physiol* 284(4): C977–87.
3. Grzeszkiewicz TM et al. (2001) CYR61 stimulates human skin fibroblast migration through Integrin alpha vbeta 5 and enhances mitogenesis through integrin alpha vbeta 3, independent of its carboxyl-terminal domain. *J Biol Chem* 276(24): 21943–50.
4. Levy L et al. (2000) beta1 integrins regulate keratinocyte adhesion and differentiation by distinct mechanisms. *Mol Biol Cell* 11(2): 453–66.
5. Juliano R. (1996) Cooperation between soluble factors and integrin-mediated cell anchorage in the control of cell growth and differentiation. *Bioessays* 18(11): 911–7.
6. Miyamoto S et al. (1995) Synergistic roles for receptor occupancy and aggregation in integrin transmembrane function. *Science* 267(5199): 883–5.
7. Schlossman S et al. (Eds.). (1995) *Leucocyte Typing V: White cell differentiation antigens*. New York: Oxford University Press.
8. Hynes RO. (1992) Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* 69(1): 11–25.
9. Defilippi P et al. (1991) Differential distribution and modulation of expression of alpha 1/beta 1 integrin on human endothelial cells. *J Cell Biol* 114(4): 855–63.
10. Sonnenberg A et al. (1990) Integrin recognition of different cell-binding fragments of laminin (P1, E3, E8) and evidence that alpha 6 beta 1 but not alpha 6 beta 4 functions as a major receptor for fragment E8. *J Cell Biol* 110(6): 2145–55.
11. Knapp W et al. (Eds.). (1989) *Leucocyte Typing IV: White cell differentiation antigens* (pp. 628–34). New York: Oxford University Press.
12. Horton MA et al. (1985) Monoclonal antibodies to osteoclastomas (giant cell bone tumors): definition of osteoclast-specific cellular antigens. *Cancer Res* 45(11): 5663–9.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists and MesenCult are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.