Anti-Human CD4 Antibody, Clone OKT4, FITC

Antibodies

Mouse monoclonal IgG2b antibody against human, rhesus, cynomolgus

CD4, FITC-conjugated

Catalog #60016FI #60016FI.1

100 Tests 5 μL/test 25 Tests 5 μL/test



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

The OKT4 antibody reacts with CD4, an ~59 kDa single-chain type 1 transmembrane glycoprotein and member of the immunoglobulin (Ig) superfamily; CD4 contains four extracellular Ig-like domains (D1 - D4). The epitope for OKT4 has been localized to the D3 domain of the protein, which has a structure resembling an Ig variable domain. CD4 is expressed at relatively high levels by most thymocytes and a subpopulation of T cells (T-helper cells), and at lower levels by peripheral blood monocytes and macrophages. CD4 binds to a non-polymorphic region of MHC II and acts as a co-receptor to the T cell receptor (TCR) in MHC II-restricted antigen recognition by enhancing the affinity of the association between the TCR and MHC II-antigen complex. CD4 also functions to amplify signals from the TCR to the cytoplasm through the interaction of its intracellular domain with cytoplasmic tyrosine kinases such as Lck. Moreover, CD4 is a receptor for human immunodeficiency virus (HIV). Binding of the OKT4 antibody to CD4 does not block HIV binding.

Target Antigen Name: CD4
Alternative Names: T4
Gene ID: 920

Species Reactivity: Human, Rhesus, Cynomolgus, Chimpanzee

Host Species: Mouse
Clonality: Monoclonal
Clone: OKT4

Isotype: IgG2b, kappa

Immunogen: Human peripheral blood T lymphocytes
Conjugate: FITC (Fluorescein isothiocyanate)

Applications

Verified: FC Reported: FC

Special Applications: This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including

EasySep™ Direct Human CD4+ T Cell Isolation Kit (Catalog #19662), EasySep™ Human CD4+ T Cell Enrichment Kit (Catalog #19052), EasySep™ Human CD3 Positive Selection Kit II (Catalog #17851), and

EasySep™ Human CD4 Positive Selection Kit II (Catalog #17852).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) bovine serum albumin

Purification: The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions. The

solution is free of unconjugated FITC.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to

light. For product expiry date, please contact techsupport@stemcell.com.

Directions for Use: For flow cytometry, the suggested use of this antibody is 5 µL per 1 x 10^6 cells in 100 µL or per 100 µL of


whole blood. It is recommended that the antibody be titrated for optimal performance for each application.


Anti-Human CD4 Antibody, Clone OKT4, FITC

Antibodies



Data





- (A) Flow cytometry analysis of human peripheral blood mononuclear cells (PBMCs) labeled with Anti-Human CD4 Antibody, Clone OKT4, FITC and Anti-Human CD45 Antibody, Clone HI30, APC (Catalog #60018AZ).
- (B) Flow cytometry analysis of human PBMCs labeled with Mouse IgG2b, kappa Isotype Control Antibody, Clone MPC-11, FITC (Catalog #60072FI) and Anti-Human CD45 Antibody, Clone HI30, APC.

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

- 1. Antiga E et al. (2014) Regulatory T cells in skin lesions and blood of patients with bullous pemphigoid. J Eur Acad Dermatol Venereol 28(2): 222–30.(FC, IF, IHC)
- 2. Buggert M et al. (2014) Multiparametric bioinformatics distinguish the CD4/CD8 ratio as a suitable laboratory predictor of combined T cell pathogenesis in HIV infection. J Immunol 192(5): 2099–108. (FC)
- 3. Kmieciak M et al. (2009) Human T cells express CD25 and Foxp3 upon activation and exhibit effector/memory phenotypes without any regulatory/suppressor function. J Transl Med 7: 89. (FC)
- 4. Rosenzweig M et al. (2001) Identification of primitive hematopoietic progenitor cells in the rhesus macague. J Med Primatol 30(1): 36–45. (FC)
- 5. Barclay AN et al. (Eds.). (1997) The Leucocyte Antigen Factsbook, Second Edition (pp. 141-2). New York: Academic Press.
- 6. Center DM et al. (1996) Interleukin 16 and its function as a CD4 ligand. Immunol Today 17(10): 476-81.
- 7. Gaubin M et al. (1996) Molecular basis of T lymphocyte CD4 antigen functions. Eur J Clin Chem Clin Biochem 34(9): 723-8.
- 8. Sakihama T et al. (1995) Molecular recognition of antigen involves lattice formation between CD4, MHC class II and TCR molecules. Immunol Today 16(12): 581–7.
- 9. Bour S et al. (1991) Inhibition of gp160 and CD4 maturation in U937 cells after both defective and productive infections by human immunodeficiency virus type 1. J Virol 65(12): 6387–96. (FC, IP)
- 10. Knapp W et al. (Eds.). (1989) Leucocyte Typing IV: White Cell Differentiation Antigens (pp. 628-34). New York: Oxford University Press.
- 11. Parnes JR. (1989) Molecular biology and function of CD4 and CD8. Adv Immunol 44: 265-311.
- 12. Linder J et al. (1987) Monoclonal antibodies marking T lymphocytes in paraffin-embedded tissue. Am J Pathol 127(1): 1-8. (IHC)
- 13. Reinherz EL et al. (1979) Separation of functional subsets of human T cells by a monoclonal antibody. Proc Natl Acad Sci USA 76(8): 4061–5. (FACS, FC)

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. Alexa Fluor is a registered trademark of Life Technologies Corporation. Antibodies conjugated to Alexa Fluor® are licensed for internal research use only and sale is expressly conditioned on the buyer not using the antibody for manufacturing, performing a service or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.