

Anti-Human CD73 Antibody, Clone AD2, PE



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Antibodies

Mouse monoclonal IgG1 antibody
against human, rhesus, chimpanzee
CD73, PE-conjugated

Catalog #60044PE	100 tests	5 µL/test
#60044PE.1	25 tests	5 µL/test

FOR RESEARCH USE ONLY. NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES.

Product Description

The AD2 antibody reacts with human CD73, a glycosyl phosphatidylinositol (GPI)-anchored glycoprotein and ecto-5'-nucleotidase expressed on the surface of subsets of B and T cells, follicular dendritic cells, mesenchymal stem cells, endothelial cells and epithelial cells. CD73 comprises a homodimer of ~70 kDa subunits that contact each other through their C-terminal domains. The enzyme catalyzes the hydrolysis of 5'-adenosine monophosphate (AMP) to form the bioactive nucleoside, adenosine, and plays a pivotal role in the activation of P1 adenosine receptors by regulating extracellular adenosine concentrations. CD73 also appears to function as a co-signaling molecule on T cells and as an adhesion molecule mediating lymphocyte interactions with the endothelium and follicular dendritic cells. CD73 is used as a marker for lymphocyte differentiation, its expression increasing during development. It is also a useful marker for identifying undifferentiated mesenchymal stem cells. CD73 is highly expressed in many types of human and mouse cancers and has been implicated in the control of tumor growth. Genetic defects in CD73 have been linked to several immunodeficiency diseases.

Target Antigen Name:	CD73
Alternative Names:	5'-nucleotidase, ecto (CD73), Ecto-5'-nucleotidase, L-VAP-2, NT5E
Gene ID:	4907
Species Reactivity:	Human, Rhesus, Chimpanzee, Pigtailed macaque
Host Species:	Mouse (BALB/c)
Clonality:	Monoclonal
Clone:	AD2
Isotype:	IgG1, kappa
Immunogen:	Human pre-B leukemia cell line 207
Conjugate:	PE

Applications


Verified:	FC
Reported:	FACS, FC, IF, IHC
Special Applications:	This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including EasySep™ Human T Cell Enrichment Kit (Catalog #19051) and EasySep™ Human B Cell Enrichment Kit (Catalog #19054), and for labeling human mesenchymal cells grown in MesenCult™-XF Medium (Catalog #05420) and MesenCult™-ACF Medium (Catalog #05440).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) bovine serum albumin
Purification:	The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com.
Directions for Use:	For flow cytometry the suggested use of this antibody is ≤ 5 µL per 1 x 10 ⁶ cells in 100 µL volume. It is recommended that the antibody be titrated for optimal performance for each application.

Data



(A) Flow cytometry analysis of human peripheral blood mononuclear cells (PBMCs; gated on lymphocytes) labeled with Anti-Human CD73 Antibody, Clone AD2, PE and Anti-Human CD3 Antibody, Clone UCHT1, Alexa Fluor® 488 (Catalog #60011AD).

(B) Flow cytometry analysis of human PBMCs (gated on lymphocytes) labeled with a mouse IgG1, kappa isotype control antibody and Anti-Human CD3 Antibody, Clone UCHT1, Alexa Fluor® 488.

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

1. Aomatsu E et al. (2014) Novel SCRG1/BST1 axis regulates self-renewal, migration, and osteogenic differentiation potential in mesenchymal stem cells. *Sci Rep* 4(3652): 1-9. (FC)
2. Allard B et al. (2014) Targeting CD73 and downstream adenosine receptor signaling in triple-negative breast cancer. *Expert Opin Ther Targets* (1): 1-19.
3. Touboul C et al. (2013) Mesenchymal stem cells enhance ovarian cancer cell infiltration through IL6 secretion in an amniochorionic membrane based 3D model. *J Transl Med* 11(28): 1-11. (FACS, FC)
4. Terp MG et al. (2013) Anti-human CD73 monoclonal antibody inhibits metastasis formation in human breast cancer by inducing clustering and internalization of CD73 expressed on the surface of cancer cells. *J Immunol* 191(8): 4165-73. (FC)
5. Hermida-Gómez T. (2011) Quantification of cells expressing mesenchymal stem cell markers in healthy and osteoarthritic synovial membranes. *J Rheumatol* 38(2): 339-49 (FC, IF, IHC)
6. Liao J et al. (2011) Cells isolated from inflamed periapical tissue express mesenchymal stem cell markers and are highly osteogenic. *J Endod* 37(9): 1217-24. (FC)
7. Tóth I et al. (2011) Decreased frequency of CD73+CD8+ T cells of HIV-infected patients correlates with immune activation and T cell exhaustion. *J Leukoc Biol* 94(4): 551-61 (FACS, FC, ICC, IF)
8. Deaglio S et al. (2007) Adenosine generation catalyzed by CD39 and CD73 expressed on regulatory T cells mediates immune suppression. *J Exp Med* 204(6): 1257-65. (FC)
9. Borrione P et al. (1999) CD38 stimulation lowers the activation threshold and enhances the alloreactivity of cord blood T cells by activating the phosphatidylinositol 3-kinase pathway and inducing CD73 expression. *J Immunol* 162(10): 6238-46. (FC)
10. Gutensohn W et al. (1995) Ecto-5'-nucleotidase activity is not required for T cell activation through CD73. *Cell Immunol* 161(2): 213-17. (FC)
11. Nakamura T et al. (1993) Characterization of an IgM Fc-binding receptor on human T cells. *J Immunol* 151(12): 6933-41. (FC)
12. Thomson LF et al. (1990) Production and characterization of monoclonal antibodies to the glycosyl phosphatidylinositol-anchored lymphocyte differentiation antigen ecto-5'-nucleotidase (CD73). *Tissue Antigens* 35(1): 9-19. (FA/Blocking, IHC, IP, WB)
13. Salazar-Gonzalez JF et al. (1985) Reduced ecto-5'-nucleotidase activity and enhanced OKT10 and HLA-DR expression on CD8 (T suppressor/cytotoxic) lymphocytes in the acquired immune deficiency syndrome: evidence of CD8 cell immaturity. *J Immunol* 135(3): 1778-85. (FA)

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485 MEDICAL DEVICE STANDARDS.

Copyright © 2015 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, EasySep, and MesenCult are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. This product is licensed for internal research use only and its sale is expressly conditioned on the buyer not using it for manufacturing, performing a service, or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.