Anti-Human OCT4 (OCT3) Antibody, Clone 3A2A20, PE

Antibodies

Mouse monoclonal IgG2b antibody against human OCT4 (OCT3), PE-

conjugated

Catalog #60093PE

100 Tests 5 µL/test

#60093PE.1 25 Tests 5 µL/test



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WERSITE

Product Description

The 3A2A20 antibody reacts with human OCT4 (octamer-binding transcription factor 4; also known as OCT3 and OCT3/4), an ~40 kDa homeodomain transcription factor belonging to the POU family, which is expressed in undifferentiated human embryonic stem (ES), induced pluripotent stem (iPS), embryonal carcinoma (EC) and embryonic germ (EG) cells. OCT4 binds to the octamer motif (5'-ATTTGCAT-3') and plays a key role in maintaining cells in a pluripotent state by interacting with other transcription factors such as SOX2 to regulate the expression of several genes, including FBX15, FGF-4, REX1, SOX2 and osteopontin. Levels of OCT4 are down-regulated during differentiation and it has thus emerged as a useful marker of pluripotency in stem cells, as well as a marker for certain human malignant germ cell tumours. Expression of OCT4 together with other transcription factors has been used to reprogram somatic cells into iPS cells. Multiple isoforms of OCT4 have been observed and in humans at least two are functionally active.

Target Antigen Name: OCT4 (OCT3)

Alternative Names: OCT-3, OCT3, OCT-4, octamer-binding transcription factor 4, POU domain class 5 transcription factor 1,

POU5F1

Gene ID: 5460 Species Reactivity: Human **Host Species:** Mouse Clonality: Monoclonal Clone: 3A2A20 Isotype: IgG2b, kappa

Immunogen: Recombinant partial human OCT4 protein (amino acids 1 - 141)

Conjugate: PF

Applications

Verified: FC Reported: FC

This antibody clone has been verified for labeling human ES and iPS cells grown in TeSR™-E8™ (Catalog Special Applications:

#05940), mTeSR™1 (Catalog #05850) and TeSR™2 (Catalog #05860).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) bovine serum albumin

Purification: The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions. The

solution is free of unconjugated PE and unconjugated antibody.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to

light. For product expiry date, please contact techsupport@stemcell.com.

Directions for Use: For flow cytometry the suggested use of this antibody is 5 µL per 1 x 10^6 cells in 100 µL volume or per

100 µL of whole blood. It is recommended that the antibody be titrated for optimal performance for each


application.


Anti-Human OCT4 (OCT3) Antibody, Clone 3A2A20, PE

Antibodies



Data





(A) Flow cytometry analysis of human ES cells cultured with mTeSR™1 on Corning® Matrigel®. The ES cells (filled histogram) or HT1080 fibrosarcoma cells (negative control; dashed line histogram) were fixed and labeled with Anti-Human OCT4 (OCT3) Antibody, Clone 3A2A20, PE. Labeling of the ES cells with Mouse IgG2b, kappa Isotype Control Antibody, Clone MPC-11, PE (Catalog #60072PE) is shown (solid line histogram).

(B) Human ES cells were cultured with TeSR™-E8™ on glass coverslips coated with Vitronectin XF™ (Catalog #07180), then fixed and labeled with

50 µm

Anti-Human OCT4 (OCT3) Antibody, Clone 3A2A20, PE. Inset shows labeling of human ES cells with Mouse IgG2b, kappa Isotype Control Antibody, Clone MPC-11, PE.

(C) DAPI counterstaining of the cells shown in figure (B); nuclear localization of the OCT4 (OCT3) marker is evident.

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

- 1. Malchenko S et al. (2014) Onset of rosette formation during spontaneous neural differentiation of hESC and hiPSC colonies. Gene 534(2): 400–7. (ICC, IF)
- 2. Pirozhkova I et al. (2013) Differences in transcription patterns between induced pluripotent stem cells produced from the same germ layer are erased upon differentiation. PLoS One 8(1): e53033. (ICC, IF)
- 3. Rijlaarsdam MA et al. (2011) Specific detection of OCT3/4 isoform A/B/B1 expression in solid (germ cell) tumours and cell lines: confirmation of OCT3/4 specificity for germ cell tumours. Br J Cancer 105(6): 854–63. (IHC)
- 4. McGuckin C et al. (2008) Culture of embryonic-like stem cells from human umbilical cord blood and onward differentiation to neural cells in vitro. Nat Protoc 3(6): 1046–55. (ICC, IF)
- 5. Takahashi K et al. (2007) Induction of pluripotent stem cells from adult human fibroblasts by defined factors. Cell 131(5): 861–72. (WB)
- 6. Wernig M et al. (2007) In vitro reprogramming of fibroblasts into a pluripotent ES-cell-like state. Nature 448(7151): 318–24. (IHC, WB)
- 7. Loh Y-H et al. (2006) The Oct4 and Nanog transcription network regulates pluripotency in mouse embryonic stem cells. Nat Genet 38(4): 431–40.
- 8. Niwa H et al. (2000) Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. Nat Genet 24(4): 372-6. (WB)
- 9. Reubinoff BE et al. (2000) Embryonic stem cell lines from human blastocysts: somatic differentiation in vitro. Nat Biotechnol 18(4): 399-404.
- 10. Nichols J et al. (1998) Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. Cell 95(3): 379–91. (ICC, IF)
- 11. Rosner MH et al. (1990) A POU-domain transcription factor in early stem cells and germ cells of the mammalian embryo. Nature 345(6277): 686–92.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists and NeuroCult are trademarks of STEMCELL Technologies Inc. Matrigel is a trademark of Corning® Incorporated. TeSR, E8 and mTeSR are trademarks of WARF. Vitronectin XF is a trademark of Primorigen Biosciences®. Vitronectin XFIII is manufactured by Primorigen Biosciences®. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. This product is licensed for internal research use only and its sale is expressly conditioned on the buyer not using it for manufacturing, performing a service, or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.