

Description

Isolate highly purified CD8+ cells from fresh or previously frozen human peripheral blood mononuclear cells (PBMCs) by immunomagnetic positive selection.

- Fast and easy-to-use
- Up to 99% purity
- · No columns required

This kit targets CD8+ cells for positive selection with an antibody recognizing the CD8 surface marker. Desired cells are labeled with antibodies and magnetic particles, and separated without columns using an EasySep[™] magnet. Unwanted cells are simply poured off, while desired cells remain in the tube. Isolated cells are immediately available for downstream applications such as flow cytometry, culture, or DNA/RNA extraction.

Component Descriptions

COMPONENT NAME	COMPONENT #	QUANTITY	STORAGE	SHELF LIFE	FORMAT
EasySep™ CD8 Positive Selection Cocktail	18053C.2	1 x 1 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A combination of monoclonal antibodies in PBS. Includes an Fc receptor blocking antibody.
EasySep™ Magnetic Nanoparticles Positive Selection	18150	1 x 1 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A suspension of magnetic particles in water.

PBS - phosphate-buffered saline

Components may be shipped at room temperature (15 - 25°C) but should be stored as indicated above.

Sample Preparation

For available fresh and frozen samples, see www.stemcell.com/primarycells.

PERIPHERAL BLOOD

Prepare a PBMC suspension from whole peripheral blood by centrifugation over a density gradient medium (e.g. Lymphoprep™, Catalog #07801). For more rapid PBMC preparation, use the SepMate™ RUO (Catalog #86450/86415) or SepMate™ IVD* (Catalog #85450/85415) cell isolation tube.

If using previously frozen PBMCs, incubate the cells with DNase I Solution (Catalog #07900) at a concentration of 100 μg/mL at room temperature (15 - 25°C) for at least 15 minutes prior to labeling and separation. Filter aggregated suspensions through a 40 μm Cell Strainer (Catalog #27305) for optimal results.

After preparation, resuspend cells at 1 x 10^8 cells/mL in recommended medium.

* SepMate[™] IVD is only available in select regions where it is registered as an In Vitro Diagnostic (IVD) device for the isolation of mononuclear cells (MNCs) from whole blood or bone marrow by density gradient centrifugation. In all other regions SepMate[™] is available for research use only (RUO).

Recommended Medium

EasySep[™] Buffer (Catalog #20144), RoboSep[™] Buffer (Catalog #20104), or PBS containing 2% fetal bovine serum (FBS) and 1 mM EDTA. Medium should be free of Ca++ and Mg++.





Directions for Use – Manual EasySep™ Protocols

See page 1 for Sample Preparation and Recommended Medium. Refer to Table 1 for detailed instructions regarding the EasySep™ procedure for each magnet.

Table 1. EasySep[™] Human CD8 Positive Selection Kit Protocol

		EASYSEP™ MAGNETS			
STEP	INSTRUCTIONS	EasySep™ (Catalog #18000)	"The Big Easy" (Catalog #18001)		
Prepare sample at the indicated cell concentration within the volume range.		1 x 10^8 cells/mL 0.1 - 2.5 mL NOTE: If starting with fewer than 1 x 10^7 cells, resuspend cells in 0.1 mL	1 x 10^8 cells/mL 0.25 - 8 mL NOTE: If starting with fewer than 2.5 x 10^7 cells, resuspend cells in 0.25 mL		
	Add sample to required tube.	5 mL (12 x 75 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352058)	14 mL (17 x 100 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352057)		
2	Add Selection Cocktail to sample.	100 μL/mL of sample	100 μL/mL of sample		
2	Mix and incubate.	RT for 15 minutes	RT for 15 minutes		
3	Mix Magnetic Particles. NOTE: Particles should appear evenly dispersed.	Pipette up and down more than 5 times	Pipette up and down more than 5 times		
4	Add Magnetic Particles to sample.	100 μL/mL of sample	100 μL/mL of sample		
4	Mix and incubate.	RT for 10 minutes	RT for 10 minutes		
5	Add recommended medium to top up the sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times.	Top up to 2.5 mL	 Top up to 5 mL for samples < 1 mL Top up to 10 mL for samples ≥ 1 mL 		
	Place the tube (without lid) into the magnet and incubate.	RT for 5 minutes	RT for 5 minutes		
6	Pick up the magnet, and in one continuous motion invert the magnet and tube,* pouring off the supernatant. Remove the tube from the magnet; this tube contains the isolated cells.	Discard supernatant	Discard supernatant		
7	Repeat steps as indicated.	Steps 5 and 6, two more times (total of 3 x 5-minute separations)	Steps 5 and 6, two more times (total of 3 x 5-minute separations)		
8	Resuspend cells in desired medium. Be sure to collect cells from the sides of the tube.	Isolated cells are ready for use	Isolated cells are ready for use		

RT - room temperature (15 - 25°C)

* Leave the magnet and tube inverted for 2 - 3 seconds, then return upright. Do not shake or blot off any drops that may remain hanging from the mouth of the tube.





Directions for Use – Fully Automated RoboSep[™] Protocol

See page 1 for Sample Preparation and Recommended Medium. Refer to Table 2 for detailed instructions regarding the RoboSep™ procedure.

Table 2. RoboSep[™] Human CD8 Positive Selection Kit Protocol

STEP	INSTRUCTIONS	RoboSep™ (Catalog #20000 and #21000)	
1	Prepare sample at the indicated cell concentration within the volume range.	1 x 10^8 cells/mL 0.25 - 8 mL NOTE: If starting with fewer than 2.5 x 10^7 cells, resuspend cells in 0.25 mL	
	Add sample to required tube.	14 mL (17 x 100 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352057)	
2	Select protocol.	 Human CD8 Positive Selection 18053-high purity Human CD8 Positive Selection 18053-high recovery 	
3	Mix Magnetic Particles. NOTE: Particles should appear evenly dispersed.	Pipette up and down more than 5 times	
4	Load the carousel.	Follow on-screen prompts	
4	Start the protocol.	Press the green "Run" button	
5	Unload the carousel when the run is complete. Remove the tube containing the isolated cells and resuspend in desired medium. Be sure to collect cells from the sides of the tube.	Isolated cells are ready for use	

Notes and Tips

ASSESSING PURITY


For purity assessment of CD8+ cells by flow cytometry use one of the following fluorochrome-conjugated antibody clones:

- Anti-Human CD8a Antibody, Clone RPA-T8 (Catalog #60022), or
- Anti-human CD8 antibody, clone HIT8a or clone B9.11, or
- Anti-Human CD8a Antibody, Clone SK1 (Catalog #60125) or clone LT8 (both partially blocked)

One of the following methods can also be used:

- Use alternative markers such as fluorochrome-conjugated Anti-Human CD3 Antibody, Clone UCHT1 (Catalog #60011) and Anti-Human CD4 Antibody, Clone OKT4 (Catalog #60016) to detect CD3+CD4- cells.
- Use a fluorochrome-conjugated secondary antibody, such as Goat Anti-Mouse IgG (H+L) Antibody, Polyclonal (Catalog #60138).

Data



Starting with fresh PBMCs, the CD8+ cell content of the isolated fraction typically ranges from 94.0 - 99.6%. In the above example, the purities of the start and final isolated fractions are 35% and 99.6%, respectively.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, EasySep, RoboSep, and SepMate are trademarks of STEMCELL Technologies Inc. Lymphoprep is a trademark of AXIS-SHIELD. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.