

Negative Selection

EasySep

Human Basophil Enrichment Kit

PROCEDURE

Version 1.0.0

CATALOG #19069



This Product Information Sheet is provided for use with RoboSep® (section A), the purple EasySep® magnet (section B) or “The Big Easy” silver EasySep® magnet (section C).

A) Fully Automated Protocol Using RoboSep® (Catalog #20000).

This procedure is used for processing 0.5 - 6.5 mL of HetaSep™-treated blood (up to 3.25×10^8 cells).

1. Prepare nucleated cell suspension (see Notes and Tips, reverse side) at a concentration of 5×10^7 cells/mL in RoboSep® Buffer (Catalog #20104). Cells must be placed in a 14 mL (17 x 100 mm) polystyrene tube to properly fit into the RoboSep® carousel.

Falcon™ 14 mL Polystyrene Round-Bottom Tubes (Becton Dickinson, Catalog #352057) are recommended.

2. Select the appropriate RoboSep® protocol:


- For most normal samples, select the protocol entitled “Human Basophil Negative Selection 19069-high purity”.

If a modified RoboSep® protocol is required, please contact StemCell Technologies' Technical Support at techsupport@stemcell.com.

3. Load the RoboSep® carousel as directed by the on-screen prompts. When all desired quadrants are loaded, press the green “Run” button. All cell labeling and separation steps will be performed by RoboSep®.

4. When cell separation is complete, collect the enriched cells in the 14 mL tube located to the left of the magnet in the second quadrant of the 2-quadrant protocol. The enriched cells are now ready for use.

Manual EasySep® Protocol Diagram



B) Manual EasySep® Protocol Using the Purple EasySep® Magnet (Catalog #18000).

This procedure is used for processing 0.5 - 2 mL of HetaSep™-treated blood (up to 1×10^8 cells).

- Prepare nucleated cell suspension at a concentration of 5×10^7 cells/mL in recommended medium (see Notes and Tips, reverse side). Cells must be placed in a 5 mL (12 x 75 mm) polystyrene tube to properly fit into the purple magnet. *Falcon™ 5 mL Polystyrene Round-Bottom Tubes (Becton Dickinson, Catalog #352058) are recommended.*
- Add EasySep® Human Basophil Enrichment Cocktail at **50 µL/mL cells** (e.g. for 2 mL of cells, add 100 µL of cocktail). Mix well and incubate at room temperature or 4°C for **10 minutes**.
- Mix EasySep® Nanoparticles to ensure that they are in a uniform suspension by vigorously pipetting more than 5 times. Vortexing is not recommended. Add the nanoparticles at **100 µL/mL cells** (e.g. for 2 mL of cells, add 200 µL of nanoparticles). Mix well and incubate at room temperature or 4°C for **10 minutes**.
- Bring the cell suspension to a **total volume** of 2.5 mL by adding recommended medium. Thoroughly mix the cells to disrupt any red blood cell aggregates by gently pipetting up and down. Place the tube (without cap) into the magnet. Set aside for **5 minutes**.
- Pick up the EasySep® Magnet, and in one continuous motion invert the magnet and tube, pouring off the desired fraction into a new 5 mL polystyrene tube. The magnetically labeled unwanted cells will remain bound inside the original tube, held by the magnetic field of the magnet. Leave the magnet and the tube in inverted position for 2 - 3 seconds, then return to upright position. **Do not shake or blot off any drops that may remain hanging from the mouth of the tube.**
- Remove the empty tube from the EasySep® magnet and place the new tube containing the supernatant fraction into the magnet. Set aside for **5 minutes**.
- Repeat Step 5. The negatively selected enriched cells in the new tube are now ready for use, or for a third round of separation, if desired (performing only 2 rounds of magnetic separation will increase cell recovery, but may reduce purity).
- Repeat Step 6, and then Step 5 once more, for a total of 3 x 5-minute separations in the magnet. The negatively selected enriched cells in the new tube are now ready for use.

C) Manual EasySep® Protocol Using “The Big Easy” Silver EasySep® Magnet (Catalog #18001).

This procedure is used for processing 0.5 - 6.5 mL of HetaSep™-treated blood (up to 3.25×10^8 cells).

- Prepare nucleated cell suspension at a concentration of 5×10^7 cells/mL in recommended medium (see Notes and Tips, reverse side). Cells must be placed in a 14 mL (17 x 100 mm) polystyrene tube to properly fit into the silver magnet. *Falcon™ 14 mL Polystyrene Round-Bottom Tubes (Catalog #352057) are recommended.*
- Add EasySep® Human Basophil Enrichment Cocktail at **50 µL/mL cells** (e.g. for 2 mL of cells, add 100 µL of cocktail). Mix well and incubate at room temperature or 4°C for **10 minutes**.
- Mix EasySep® Nanoparticles to ensure that they are in a uniform suspension by pipetting vigorously more than 5 times. Vortexing is not recommended. Add the particles at **100 µL/mL cells** (e.g. for 2 mL of cells, add 200 µL of nanoparticles). Mix well and incubate at room temperature or 4°C for **10 minutes**.
- Bring the cell suspension to a **total volume** of 5.0 mL (for $<10^8$ cells) or 10 mL (for $1 - 3.25 \times 10^8$ cells) by adding recommended medium. Thoroughly mix the cells to disrupt any red blood cell aggregates by gently pipetting up and down. Place the tube (without cap) into the magnet. Set aside for **10 minutes**.
- Pick up the EasySep® Magnet, and in one continuous motion invert the magnet and tube, pouring off the desired fraction into a new 14 mL polystyrene tube. The magnetically labeled unwanted cells will remain bound inside the original tube, held by the magnetic field of the magnet. Leave the magnet and the tube in inverted position for 2 - 3 seconds, then return to upright position. **Do not shake or blot off any drops that may remain hanging from the mouth of the tube.**
- Remove the empty tube from the EasySep® magnet and place the new tube containing the supernatant fraction into the magnet. Set aside for **10 minutes**.
- Repeat Step 5 for a total of 2 x 10-minute separations in the magnet. The negatively selected enriched cells in the new tube are now ready for use.

StemCell Technologies

In North America
Tel: 1.604.877.0713
Fax: 1.604.877.0704
Toll Free Tel: 1.800.667.0322
Toll Free Fax: 1.800.567.2899
e-mail: info@stemcell.com
www.stemcell.com

In the United Kingdom
Tel: +44.(0).20.7691.3561
Fax: +33.(0).4.76.18.99.63
Toll Free within United Kingdom:
Tel: 0800.731.27.14
Fax: 0800.731.27.13
e-mail: info@stemcellgb.com

In Europe
Tel: +33.(0).4.76.04.75.30
Fax: +33.(0).4.76.18.99.63
e-mail: info@stemcellfrance.com

June 2007

FOR RESEARCH USE ONLY

#29109

Catalog #19069

For labeling 10^9 total cells

Components:

- EasySep[®] Human Basophil Enrichment Cocktail 1.0 mL
- EasySep[®] Magnetic Nanoparticles 3 x 1.0 mL



Product Information Sheet

REQUIRED EQUIPMENT:

EasySep[®] Magnet (Catalog #18000), "The Big Easy" EasySep[®] Magnet (Catalog #18001), or RoboSep[®] (Catalog #20000).

PRODUCT DESCRIPTION AND APPLICATIONS:

The EasySep[®] Human Basophil Enrichment Kit is designed to enrich human basophils by depleting non-basophils. Start cell samples are prepared by performing a HetaSep[™] sedimentation of the peripheral blood (see Notes and Tips, below).

EASYSEP[®] LABELING OF HUMAN CELLS:

Unwanted cells are specifically labeled with dextran-coated magnetic nanoparticles using bispecific Tetrameric Antibody Complexes (TAC). These complexes recognize both dextran and the cell surface antigen expressed on the unwanted cells (Figure 1). The small size of the magnetic dextran iron particles allows for efficient binding to the TAC-labeled cells. Magnetically labeled cells are then separated from unlabeled target cells using the EasySep[®] procedure (reverse side).




Figure 1.

Schematic Drawing of EasySep[®] TAC Magnetic Labeling of Human Cells.

NOTES AND TIPS:

Preparing a Nucleated Cell Suspension for Isolation of Basophils.

Collect whole blood in a blood collection tube containing heparin or other anticoagulant. Add 1 part HetaSep[™] (Catalog #07906) to 5 parts blood and mix well. Use the minimum sized tube for the total volume of HetaSep[™]: blood sample. Centrifuge for 6 minutes at $110 \times g$ at room temperature with the brake off. Remove tube from centrifuge and let sit until the red blood cell : plasma interface is at approximately 40% of the total volume (a maximum of 10 minutes). Carefully remove all of the plasma containing the nucleated cells (everything above the red blood cell pellet) and wash this fraction, using 4 parts cold recommended medium to 1 part recovered plasma / cells. Centrifuge for 10 minutes at $500 \times g$ with the brake set to low. Discard supernatant and wash a second time to remove excess platelets, centrifuging for 10 minutes at $120 \times g$ with the brake off. Discard supernatant and resuspend at 5×10^7 cells/mL.

Do not use dextran sedimentation to prepare cells.

Optimal Cell Number. We do not recommend using fewer than 5×10^7 cells per separation as this may result in sub-optimal performance.



Recommended Medium. The recommended medium is RoboSep[®] Buffer (Catalog #20104), or Phosphate Buffered Saline (PBS) containing 2% FBS (Catalog #07905) and 1 mM EDTA. Media should be Ca^{++} and Mg^{++} free.

Assessing Purity. Purity of basophils can be measured by flow cytometry after simultaneously staining with fluorochrome-conjugated anti-CD123 (e.g. anti-CD123 PE, Catalog #10532), and anti-IgE or anti-CD203c. Basophils are defined as $CD123^+IgE^+$ or $CD123^+CD203c^{lo}$. Alternatively, purity may be assessed by performing a cytospin on the enriched cells followed by Wright's or May-Grünwald staining (e.g. Sigma-Aldrich Catalog #W0625 or #205435).

TYPICAL EASYSEP[®] BASOPHIL ENRICHMENT PROFILE:

Start: $1.5\% CD123^+IgE^+$

Enriched: $98\% CD123^+IgE^+$



Starting with freshly prepared nucleated cells, the basophil content of the enriched fraction typically ranges from 92 - 99% (gated on $CD45^+$).

COMPONENT DESCRIPTIONS:

EasySep[®] Negative Selection Human Basophil Enrichment Cocktail code #19069C

This cocktail contains a combination of monoclonal antibodies purified from hybridoma culture supernatant by affinity chromatography using Protein A or Protein G Sepharose. These antibodies are bound in bispecific Tetrameric Antibody Complexes (TAC) which are directed against cell surface antigens on human blood cells (CD2, CD3, CD14, CD15, CD16, CD19, CD24, CD34, CD36, CD45RA, CD56, glycoprotein A) and dextran. The mouse monoclonal antibody subclass is IgG_1 . It should be noted that this product is a biological reagent, and as such cannot be completely characterized or quantified. Some variability is unavoidable.

EasySep[®] Magnetic Nanoparticles code #19150.1

A suspension of magnetic dextran iron particles in water.

STABILITY AND STORAGE:

EasySep[®] Human Basophil Enrichment Cocktail

Stable at $4^\circ C$ for 2 years. Do not freeze this product. Contents sterile in unopened tube. This product may be shipped at room temperature, and should be refrigerated upon receipt.

EasySep[®] Magnetic Nanoparticles

Stable at $4^\circ C$ for 2 years. Contents sterile in unopened tube. This product may be shipped at room temperature, and should be refrigerated upon receipt.

StemCell Technologies

In North America

Tel: 1.604.877.0713
Fax: 1.604.877.0704
Toll Free Tel: 1.800.667.0322
Toll Free Fax: 1.800.567.2899
e-mail: info@stemcell.com
www.stemcell.com

In the United Kingdom

Tel: +44.(0).20.7691.3561
Fax: +33.(0).4.76.18.99.63
Toll Free within United Kingdom: e-mail: info@stemcellfrance.com
Tel: 0800.731.27.14
Fax: 0800.731.27.13
e-mail: info@stemcellgb.com

In Europe

Tel: +33.(0).4.76.04.75.30
Fax: +33.(0).4.76.18.99.63

FOR RESEARCH USE ONLY

#29109