



EasySep™ Human Pan-Granulocyte Isolation Kit

Negative Selection
Catalog #19259

For processing 1 x 10⁹ cells



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713
INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM
FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Document #29297 | Version 2_0_0

Description

Isolate highly purified pan-granulocytes (neutrophils, eosinophils, and basophils) from fresh human peripheral blood leukocytes by immunomagnetic negative selection.

- Fast, easy-to-use and column-free
- Up to 99% purity
- Untouched, viable cells

This kit targets non-granulocytes for removal with antibodies recognizing specific cell surface markers. Unwanted cells are labeled with antibodies and magnetic particles, and separated without columns using an EasySep™ magnet. Desired cells are simply poured off into a new tube. Isolated cells are immediately available for downstream applications such as flow cytometry, culture, or DNA/RNA extraction.

Component Descriptions

COMPONENT NAME	COMPONENT #	QUANTITY	STORAGE	SHELF LIFE	FORMAT
EasySep™ Human Pan-Granulocyte Isolation Cocktail	19259C	1 x 1 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A combination of monoclonal antibodies in PBS.
EasySep™ D2 Magnetic Particles	19650	2 x 1 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A suspension of magnetic particles in TBS.

PBS - phosphate-buffered saline; TBS - TRIS-buffered saline

Components may be shipped at room temperature (15 - 25°C) but should be stored as indicated above.

Sample Preparation

Important: Do not use dextran sedimentation to prepare cells.

WHOLE BLOOD USING RED BLOOD CELL (RBC) LYSIS (preferred for slightly higher recovery)

1. Collect whole blood in a blood collection tube containing anticoagulant.
2. Add 4 parts Ammonium Chloride Solution (Catalog #07800) to 1 part whole blood and mix well.
3. Incubate on ice for 15 minutes then centrifuge at 300 x g for 10 minutes.
4. Wash pellet once with recommended medium and centrifuge at 120 x g for 10 minutes with the brake off.
5. Discard supernatant and resuspend cells at 5 x 10⁷ cells/mL in recommended medium.

WHOLE BLOOD USING HETASEP™ RBC SEDIMENTATION (preferred for faster, lysis-free sample processing)

1. Collect whole blood in a blood collection tube containing anticoagulant.
2. Add 1 part HetaSep™ (Catalog #07906) to 5 parts whole blood and mix well. Use the minimum-sized tube for the total volume of HetaSep™:blood sample. A 14 mL tube is the maximum size recommended for optimal leukocyte recovery.
3. Centrifuge sample at 90 x g for 2 minutes (if total volume ≤ 5 mL) or 5 minutes (if total volume > 5 mL) at room temperature (15 - 25°C) with the brake off.
4. Remove tube from centrifuge and let sit undisturbed for 10 minutes.
5. Harvest the leukocyte-rich plasma (everything above the RBC fraction) into a 50 mL tube and add 4 parts recommended medium to 1 part harvested cells/plasma.
6. Centrifuge at 300 x g for 10 minutes at room temperature.
7. Discard supernatant and wash pellet to remove excess platelets, centrifuging at 120 x g for 10 minutes at room temperature with the brake off.
8. Discard supernatant and resuspend cells at 5 x 10⁷ cells/mL in recommended medium.



Recommended Medium

EasySep™ Buffer (Catalog #20144), RoboSep™ Buffer (Catalog #20104), or PBS containing 2% fetal bovine serum (FBS) and 1 mM EDTA. Medium should be free of Ca⁺⁺ and Mg⁺⁺.

Directions for Use – Manual EasySep™ Protocols

See page 1 for Sample Preparation and Recommended Medium. Refer to Tables 1 and 2 for detailed instructions regarding the EasySep™ procedure for each magnet.


Table 1. EasySep™ Human Pan-Granulocyte Isolation Kit Protocol

		EASYSEP™ MAGNETS	
STEP	INSTRUCTIONS	 EasySep™ (Catalog #18000)	“The Big Easy” (Catalog #18001) 
1	Prepare sample at the indicated cell concentration within the volume range.	5 x 10 ⁷ cells/mL 0.25 - 2 mL	5 x 10 ⁷ cells/mL 0.5 - 8 mL
	Add sample to required tube.	5 mL (12 x 75 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352058)	14 mL (17 x 100 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352057)
2	Add Isolation Cocktail to sample.	50 µL/mL of sample	50 µL/mL of sample
	Mix and incubate.	RT for 10 minutes	RT for 10 minutes
3	Vortex Magnetic Particles. NOTE: Particles should appear evenly dispersed.	30 seconds	30 seconds
4	Add Magnetic Particles to sample.	50 µL/mL of sample	50 µL/mL of sample
	Mix and incubate.	RT for 5 minutes	RT for 5 minutes
5	Add recommended medium to top up the sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times.	Top up to 2.5 mL	<ul style="list-style-type: none"> • Top up to 5 mL for samples ≤ 4 mL • Top up to 10 mL for samples > 4 mL
	Place the tube (without lid) into the magnet and incubate.	RT for 5 minutes	RT for 5 minutes
6	Pick up the magnet, and in one continuous motion invert the magnet and tube,* pouring the enriched cell suspension into a new tube.	Isolated cells are ready for use	Isolated cells are ready for use

RT - room temperature (15 - 25°C)

* Leave the magnet and tube inverted for 2 - 3 seconds, then return upright. Do not shake or blot off any drops that may remain hanging from the mouth of the tube.

Table 2. EasySep™ Human Pan-Granulocyte Isolation Kit Protocol

		EASYSEP™ MAGNETS	
STEP	INSTRUCTIONS	Easy 50 (Catalog #18002)	
1	Prepare sample at the indicated cell concentration within the volume range.	5 x 10 ⁷ cells/mL 1 - 40 mL	
	Add sample to required tube.	50 mL conical tube (e.g. Corning Catalog #352070)	
2	Add Isolation Cocktail to sample.	50 µL/mL of sample	
	Mix and incubate.	RT for 10 minutes	
3	Vortex Magnetic Particles. NOTE: Particles should appear evenly dispersed.	30 seconds	
4	Add Magnetic Particles to sample.	75 µL/mL of sample	
	Mix and incubate.	RT for 10 minutes	
5	Add recommended medium to top up sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times.	<ul style="list-style-type: none"> Top up to 10 mL for samples < 5 mL Top up to 25 mL for samples ≥ 5 - 15 mL Top up to 50 mL for samples > 15 - 40 mL 	
	Place the tube (without lid) into the magnet and incubate.	RT for 10 minutes	
6	Carefully pipette** (do not pour) the enriched cell suspension into a new tube.	Use a new 50 mL tube	
7	Remove the tube from the magnet and place the new tube (without lid) into the magnet and incubate for a second separation.	RT for 5 minutes	
8	Carefully pipette** (do not pour) the enriched cell suspension into a new tube.	Isolated cells are ready for use	

** Collect the entire supernatant, all at once, into a single pipette (e.g. for volumes of 5 - 10 mL use a 10 mL serological pipette, for volumes > 10 - 25 mL use a 25 mL serological pipette, and for volumes > 25 mL use a 50 mL serological pipette).

Directions for Use – Fully Automated RoboSep™ Protocol

See page 1 for Sample Preparation and Recommended Medium. Refer to Table 3 for detailed instructions regarding the RoboSep™ procedure.

Table 3. RoboSep™ Human Pan-Granulocyte Isolation Kit Protocol

STEP	INSTRUCTIONS	RoboSep™ (Catalog #20000 and #21000)
1	Prepare sample at the indicated cell concentration within the volume range.	5 x 10 ⁷ cells/mL 0.5 - 8 mL
	Add sample to required tube.	14 mL (17 x 100 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352057)
2	Select protocol.	Human Pan-Granulocyte Isolation 19259
3	Vortex Magnetic Particles. NOTE: Particles should appear evenly dispersed.	30 seconds
4	Load the carousel.	Follow on-screen prompts
	Start the protocol.	Press the green "Run" button
5	Unload the carousel when the run is complete.	Isolated cells are ready for use

Notes and Tips

ASSESSING PURITY


For purity assessment of pan-granulocytes (neutrophils, eosinophils, and basophils) by flow cytometry use the following fluorochrome-conjugated antibody clones:

- Anti-Human CD16 Antibody, Clone 3G8 (Catalog #60041), and
- Anti-Human CD66b Antibody, Clone G10F5 (Catalog #60086), and
- Anti-Human CD123 Antibody (IL-3R α), Clone 6H6 (Catalog #60110; optional, for basophil assessment), and
- Anti-human IgE antibody (optional, for basophil assessment), and
- Anti-human CD203c antibody (optional, for basophil assessment)

Neutrophils are CD66b+CD16+, eosinophils are CD66b+CD16- and low in forward scatter but high in side scatter, and basophils are CD66-CD123+IgE+ or CD66-CD123+CD203c^{low}.

Alternatively, purity may be assessed by performing a cytospin of the enriched cells followed by Wright's or May-Grünwald staining (e.g. Sigma-Aldrich Catalog #W0625 or #205435, respectively).

Data



Starting with whole peripheral blood, the total granulocyte content of the isolated fraction typically ranges from 97 - 99%. In the above example, neutrophils are typically CD66b+CD16+, eosinophils are typically CD66b+CD16-, and basophils are CD66b- and can be further defined as CD203c+CD123+ and IgE+ (not shown).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, EasySep, HetaSep, and RoboSep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.