Antibodies	Anti-Mouse CD117 Antibody, Clone 2B8, APC		STENCELL ^M
	Rat monc against m conjugate	oclonal IgG2b antibody nouse CD117 (c-Kit), APC- ed	Scientists Helping Scientists™ │ WWW.STEMCELL.COM
Catalog #60025AZ #60025AZ.1	100 μg 25 μg	0.2 mg/mL 0.2 mg/mL	INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

The 2B8 antibody reacts with CD117 (c-Kit), an ~145 kDa type 1 transmembrane receptor for c-Kit ligand (stem cell factor/steel factor) that is broadly expressed on hematopoietic stem cells in bone marrow, including pluripotent and erythroid progenitor cells and B and T lymphocyte precursors, as well as on mast cells. CD117 belongs to the tyrosine kinase receptor family and possesses five immunoglobulin-like C2-type domains and a cytoplasmic protein kinase domain. Binding of c-Kit ligand to CD117 induces dimerization and autophosphorylation, which activates several intracellular signaling pathways critical for the proliferation and differentiation of hematopoietic stem cells. Signaling by CD117 is modulated by phosphatases and by rapid endocytosis and degradation of the receptor. The 2B8 antibody does not block binding of c-Kit ligand. Mutations in CD117 are associated with various types of tumors and the piebald trait, an autosomal dominant abnormality of pigmentation.

Target Antigen Name:	CD117 (c-Kit)
Alternative Names:	c-KIT, cKIT, Stem cell factor receptor (SCFR)
Gene ID:	16590
Species Reactivity:	Mouse
Host Species:	Rat (Wistar)
Clonality:	Monoclonal
Clone:	2B8
Isotype:	lgG2b, kappa
Immunogen:	Mouse bone marrow mast cells
Conjugate:	APC

Applications

Verified:	FC
Reported:	FC
Special Applications:	This antibody clone has been verified for purity assessments of cells isolated with EasySep [™] kits, including EasySep [™] Mouse Hematopoietic Progenitor Cell Isolation Kit (Catalog #19856).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Purification:	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com.
Directions for Use:	For flow cytometry the suggested use of this antibody is $\leq 1 \ \mu$ g per 1 x 10^6 cells in 100 μ L volume. It is recommended that the antibody be titrated for optimal performance for each application.



Data



(A) Flow cytometry analysis of C57BL/6 mouse bone marrow cells labeled with Anti-Mouse CD117 Antibody, Clone 2B8, APC and Anti-Mouse CD45R Antibody, Clone RA3-6B2, FITC (Catalog #60019FI).

(B) Flow cytometry analysis of C57BL/6 mouse bone marrow cells labeled with Rat IgG2b, kappa Isotype Control Antibody, Clone RTK4530, APC (Catalog #60077AZ) and Anti-Mouse CD45R Antibody, Clone RA3-6B2, FITC.

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

1. Gough DJ et al. (2014) STAT3 supports experimental K-RasG12D-induced murine myeloproliferative neoplasms dependent on serine phosphorylation. Blood 124(14): 2252–61. (FC)

2. Reeh KAG et al. (2014) Ectopic TBX1 suppresses thymic epithelial cell differentiation and proliferation during thymus organogenesis. Development 141(15): 2950–8. (FC)

3. Sheng KC et al. (2014) IL-3 and CSF-1 interact to promote generation of CD11c+ IL-10-producing macrophages. PLoS One 9(4): e95208.(FC)

4. Charles N et al. (2010) Basophils and the T helper 2 environment can promote the development of lupus nephritis. Nat Med 16(6): 701–7. (FC)

5. Podd BS et al. (2006) T cells in cryptopatch aggregates share TCR gamma variable region junctional sequences with gamma delta T cells in the small intestinal epithelium of mice. J Immunol 176(11): 6532–42. (IHC)

6. Wognum AW et al. (2003) Identification and isolation of hematopoietic stem cells. Arch Med Res 34(6): 461–75. (CellSep)

7. Pulendran B et al. (1997) Developmental pathways of dendritic cells in vivo: distinct function, phenotype, and localization of dendritic cell subsets in FLT3 ligand-treated mice. J Immunol 159(5): 2222–31. (FC)

8. Rico-Vargas SA et al. (1994) c-kit expression by B cell precursors in mouse bone marrow. Stimulation of B cell genesis by in vivo treatment with anti-ckit antibody. J Immunol 152(6): 2845–52. (FA/Blocking, ICC, IF)

9. Fleming WH et al. (1993) Steel factor influences the distribution and activity of murine hematopoietic stem cells in vivo. Proc Natl Acad Sci USA 90(8): 3760–4. (FC)

10. Matsuzaki Y et al. (1993) Characterization of c-kit positive intrathymic stem cells that are restricted to lymphoid differentiation. J Exp Med 178(4): 1283–92. (FACS, FC)

11. Ikuta K & Weissman IL. (1992) Evidence that hematopoietic stem cells express mouse c-kit but do not depend on steel factor for their generation. Proc Natl Acad Sci USA 89(4): 1502–6. (FC, IP)

12. Ogawa M et al. (1991) Expression and function of c-kit in hemopoietic progenitor cells. J Exp Med 174(1): 63–71. (FA, FC)

13. Flanagan JG & Leder P. (1990) The kit ligand: a cell surface molecule altered in steel mutant fibroblasts. Cell 63(1): 185–94.

14. Chabot B et al. (1988) The proto-oncogene c-kit encoding a transmembrane tyrosine kinase receptor maps to the mouse W locus. Nature 335(6185): 88–9.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists and EasySep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. This product is licensed for internal research use only and its sale is expressly conditioned on the buyer not using it for manufacturing, performing a service, or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.