

## Anti-Mouse TER119 Antibody, Clone TER-119, APC



Scientists Helping Scientists™ | [WWW.STEMCELL.COM](http://WWW.STEMCELL.COM)

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

[INFO@STEMCELL.COM](mailto:INFO@STEMCELL.COM) • [TECHSUPPORT@STEMCELL.COM](mailto:TECHSUPPORT@STEMCELL.COM)

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## Antibodies

Rat monoclonal IgG2b antibody  
against mouse TER119, APC-  
conjugated

Catalog #60033AZ  
#60033AZ.1

100 µg 0.2 mg/mL  
25 µg 0.2 mg/mL

## Product Description

The TER-119 antibody reacts with murine TER119 (Ly-76), an ~52 kDa protein associated with glycophorin A on the surface of cells of the erythroid lineage in embryonic yolk sac, fetal and newborn liver, adult bone marrow, peripheral blood, and lymphoid organs. TER119 is an erythroid-specific marker expressed at all stages of differentiation from early proerythroblasts to mature erythrocytes, but not by erythroid colony-forming cells (BFU-E, blast-forming unit erythroid, or CFU-E, colony-forming unit erythroid). The TER-119 antibody is a component of the "lineage cocktail" used to detect, or deplete cells committed to hematopoietic lineages. In adult mice, TER119 is found on ~20 - 25% of bone marrow cells and ~2 - 3% of splenocytes.

Target Antigen Name:	TER119
Alternative Names:	Ly-76, TER-119
Gene ID:	104231
Species Reactivity:	Mouse
Host Species:	Rat (WI)
Clonality:	Monoclonal
Clone:	TER-119
Isotype:	IgG2b, kappa
Immunogen:	Mouse (C57BL/6) fetal liver cells
Conjugate:	APC

## Applications


Verified:	FC
Reported:	FC
Special Applications:	This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including EasySep™ Mouse CD4+ T Cell Isolation Kit (Catalog #19852).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

## Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Purification:	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. For product expiry date, please contact <a href="mailto:techsupport@stemcell.com">techsupport@stemcell.com</a> .
Directions for Use:	For flow cytometry the suggested use of this antibody is $\leq 1 \mu\text{g}$ per $1 \times 10^6$ cells in 100 µL volume. It is recommended that the antibody be titrated for optimal performance for each application.

## Data



Flow cytometry analysis of C57BL/6 mouse bone marrow cells labeled with Anti-Mouse TER119 Antibody, Clone TER-119, APC (filled histogram) or Rat IgG2b, kappa Isotype Control Antibody, Clone RTK4530, APC (Catalog #60077AZ) (solid line histogram).

## Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at [www.stemcell.com/antibodies](http://www.stemcell.com/antibodies) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## References

1. Cui S et al. (2015) Compound loss of function of nuclear receptors Tr2 and Tr4 leads to induction of murine embryonic  $\beta$ -type globin genes. *Blood* 125(9): 1477–87. (FACS, FC)
2. Suenaga F et al. (2015) Loss of lymph node fibroblastic reticular cells and high endothelial cells is associated with humoral immunodeficiency in mouse graft-versus-host disease. *J Immunol* 194(1): 398–406. (FC)
3. Rehn M et al. (2014) Hypoxic induction of vascular endothelial growth factor regulates erythropoiesis but not hematopoietic stem cell function in the fetal liver. *Exp Hematol* 42(11): 941–4.e1. (FC)
4. Morioka S et al. (2012) TAK1 kinase signaling regulates embryonic angiogenesis by modulating endothelial cell survival and migration. *Blood* 120(18): 3846–57.
5. Sung JH et al. (2008) Isolation and characterization of mouse mesenchymal stem cells. *Transplant Proc* 40(8): 2649–54. (IHC, FC)
6. Chappaz S et al. (2007) Increased TSLP availability restores T- and B-cell compartments in adult IL-7 deficient mice. *Blood* 110(12): 3862–70. (FC)
7. Heuser M et al. (2007) MN1 overexpression induces acute myeloid leukemia in mice and predicts ATRA resistance in patients with AML. *Blood* 110(5): 1639–47. (FC)
8. Grisendi S et al. (2005) Role of nucleophosmin in embryonic development and tumorigenesis. *Nature* 437(7055): 147–53. (FC)
9. Kina T et al. (2000) The monoclonal antibody TER-119 recognizes a molecule associated with glycophorin A and specifically marks the late stages of murine erythroid lineage. *Br J Haematol* 109(2): 280–87. (IP, WB)
10. Vannucchi AM et al. (2000) Identification and characterization of a bipotent (erythroid and megakaryocytic) cell precursor from the spleen of phenylhydrazinetreated mice. *Blood* 95(8): 2559–68.
11. Kitajima K et al. (1999) Definitive but not primitive hematopoiesis is impaired in jumonji mutant mice. *Blood* 93(1): 87–95. (IHC)
12. Maraskovsky E et al. (1996) Dramatic increase in the numbers of functionally mature dendritic cells in Flt3 ligand-treated mice: multiple dendritic cell subpopulations identified. *J Exp Med* 184(5): 1953–62. (FA)
13. Ikuta K et al. (1990) A developmental switch in thymic lymphocyte maturation potential occurs at the level of hematopoietic stem cells. *Cell* 62(5): 863–74. (FC, IP, WB)

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. This product is licensed for internal research use only and its sale is expressly conditioned on the buyer not using it for manufacturing, performing a service, or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or [outlicensing@lifetech.com](mailto:outlicensing@lifetech.com). While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.