

## Anti-Human CD83 Antibody, Clone HB15e, APC



Scientists Helping Scientists™ | [WWW.STEMCELL.COM](http://WWW.STEMCELL.COM)

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

[INFO@STEMCELL.COM](mailto:INFO@STEMCELL.COM) • [TECHSUPPORT@STEMCELL.COM](mailto:TECHSUPPORT@STEMCELL.COM)

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## Antibodies

Mouse monoclonal IgG1 antibody  
against human, rhesus, cynomolgus  
CD83, APC-conjugated

Catalog #60107AZ  
#60107AZ.1

100 Tests 5 µL/test  
25 Tests 5 µL/test

## Product Description

The HB15e antibody reacts with CD83, an ~45 kDa type I transmembrane glycoprotein and member of the Siglec (sialic acid-binding immunoglobulin-like lectin) protein family, containing an extracellular V-type Ig-like domain and a C-terminal cytoplasmic tail. A soluble form of CD83 has also been identified. CD83 is expressed predominantly on circulating and tissue dendritic cells, including Langerhan cells and interdigitating reticulum cells. It is expressed weakly on activated lymphocytes and may also be found on neutrophils, monocytes and macrophages. Expression of CD83 is typically transient, except on dendritic cells, and dependent on the cell's state or maturation stage. CD83 expression correlates with upregulation of HLA class II antigen and serves as a marker for dendritic cell maturation and activation. Its function, though ill-defined, is essential for the development and maintenance of the immune system and evidence suggests it may be involved in cell adhesion and signaling, antigen presentation, and activation and proliferation of lymphocytes. The soluble form of CD83 appears to be immunosuppressive, blocking T cell activation and inhibiting dendritic cell maturation. The epitope on CD83 bound by the HB15e antibody reportedly overlaps with that of antibody 1E11.

Target Antigen Name:	CD83
Alternative Names:	B-cell activation protein, BL11, HB15
Gene ID:	9308
Species Reactivity:	Human, Rhesus, Cynomolgus, Baboon, Chimpanzee, Pigtailed Macaque
Host Species:	Mouse
Clonality:	Monoclonal
Clone:	HB15e
Isotype:	IgG1, kappa
Immunogen:	Human CD83-transfected African green monkey COS-7 cells
Conjugate:	APC

## Applications


Verified:	FC
Reported:	FC
Special Applications:	This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including EasySep™ Human CD14 Positive Selection Kit (Catalog #18058), and for analyzing human monocyte-derived mature dendritic cells cultured using StemSpan™-ACF medium (Catalog #09805).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

## Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) bovine serum albumin
Purification:	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. For product expiry date, please contact <a href="mailto:techsupport@stemcell.com">techsupport@stemcell.com</a> .
Directions for Use:	For flow cytometry the suggested use of this antibody is ≤ 5 µL per 1 × 10 <sup>6</sup> cells in 100 µL volume or 5 µL per 100 µL of whole blood. It is recommended that the antibody be titrated for optimal performance for each application.

## Data



Flow cytometry analysis of dendritic cells derived from human peripheral blood mononuclear cells (PBMCs). Cells were processed with the EasySep™ Human CD14 Positive Selection Kit and labeled with Anti-Human CD83 Antibody, Clone HB15e, APC (filled histogram) or Mouse IgG1, kappa Isotype Control Antibody, Clone MOPC-21, APC (Catalog #60070AZ) (solid line histogram).

## Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at [www.stemcell.com/antibodies](http://www.stemcell.com/antibodies) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## References

1. Pelletier S et al. (2013) Sustained hyperresponsiveness of dendritic cells is associated with spontaneous resolution of acute hepatitis C. *J Virol* 87(12): 6769–81. (FC)
2. Séguier S et al. (2013) Inhibition of the differentiation of monocyte-derived dendritic cells by human gingival fibroblasts. *PLoS One* 8(8): e70937. (FC)
3. Denniston AK et al. (2012) Aqueous humor suppression of dendritic cell function helps maintain immune regulation in the eye during human uveitis. *Invest Ophthalmol Vis Sci* 53(2): 888–96. (FC)
4. Yin Z et al. (2012) Type III IFNs are produced by and stimulate human plasmacytoid dendritic cells. *J Immunol* 189(6): 2735–45. (FC)
5. Chen L et al. (2011) Two novel monoclonal antibodies produced against human CD83 molecule. *Hybridoma (Larchmt)* 30(3): 297–302. (FC, Immunoblotting)
6. Denniston AK et al. (2011) Endogenous cortisol and TGF-beta in human aqueous humor contribute to ocular immune privilege by regulating dendritic cell function. *J Immunol* 186(1): 305–11. (FC)
7. Hovden A-O et al. (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. *BMC Immunol* 12: 2. (FC)
8. Cao W et al. (2005) CD83 is preformed inside monocytes, macrophages and dendritic cells, but it is only stably expressed on activated dendritic cells. *Biochem J* 385(1): 85–93. (FC, ICC, IF, WB)
9. Iking-Konert C et al. (2002) Up-regulation of the dendritic cell marker CD83 on polymorphonuclear neutrophils (PMN): divergent expression in acute bacterial infections and chronic inflammatory disease. *Clin Exp Immunol* 130(3): 501–8. (FC, IP)
10. Loré K et al. (2002) Accumulation of DC-SIGN+CD40+ dendritic cells with reduced CD80 and CD86 expression in lymphoid tissue during acute HIV-1 infection. *AIDS* 16(5): 683–92. (ICH)

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, EasySep and StemSpan are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. This product is licensed for internal research use only and its sale is expressly conditioned on the buyer not using it for manufacturing, performing a service, or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or [outlicensing@lifetech.com](mailto:outlicensing@lifetech.com). While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.