
Small Molecules	UM729	STENCELL™ T E C H N O L O G I E S
	5 Pyrimido-indole derivative that enhances HSC self-renewal in vitro	Scientists Helping Scientists [™] WWW.STEMCELL.COM
Catalog # 72332		TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713
	250 μg of active co	INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM
72334	1 mg of active com	FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

UM729 is a pyrimido-[4,5-b]-indole derivative which enhances the self-renewal of human hematopoietic stem cells (HSCs) in vitro (Fares et al.). UM729 does not inhibit the aryl hydrocarbon receptor (AhR) pathway, but has been shown to collaborate with AhR antagonists in preventing differentiation of acute myeloid leukemia (AML) cells in culture (Pabst et al.).

Molecular Name:	UM729
Alternative Names:	UM-729
CAS Number:	Not applicable
Chemical Formula:	$C_{20}H_{25}N_5O_2 \cdot X HCI [X H2O]$
Molecular Weight:	367.4 g/mol (free base)
Purity:	≥ 95%
Chemical Name:	Methyl 4-((3-(piperidin-1-yl)propyl)amino)-9H-pyrimido[4,5-b] indole-7-carboxylate
Structure:	
	A H



Properties

-	
Physical Appearance:	Light yellow or green to off-white solid
Storage:	Product stable at room temperature (15 - 25°C) as supplied. Protect from prolonged exposure to light. Stable as supplied for 12 months from date of receipt.
Solubility:	\cdot DMSO \leq 20 mM
	For example, to prepare a 5 mM stock solution in DMSO, dissolve 1 mg in 544 μ L of fresh DMSO or dissolve 250 μ g in 136 μ L of fresh DMSO.
	Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.
	Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to

be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.



Published Applications

MAINTENANCE AND SELF-RENEWAL

· Enhances human hematopoietic stem cell self-renewal in vitro (Fares et al.).

CANCER RESEARCH

· Collaborates with StemRegenin 1 (SR1; Catalog #72342) in preventing differentiation of AML cells in culture (Pabst et al.).

References

Fares I et al. (2014) Pyrimidoindole derivatives are agonists of human hematopoietic stem cell self-renewal. Science 345(6203): 1509–12. Pabst C et al. (2014) Identification of small molecules that support human leukemia stem cell activity ex vivo. Nat Methods 11(4): 436–42.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, visit www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is potentially hazardous. Please refer to the Safety Data Sheet (SDS).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.