

Small Molecules

16,16-Dimethyl Prostaglandin E2

Prostanoid pathway activator; Inhibits 15-hydroxy PGDH

Catalog # 72372

5 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

16,16-Dimethyl Prostaglandin E2 (16,16-Dimethyl PGE2), a stabilized derivative of PGE2, is a competitive inhibitor of 15-hydroxy prostaglandin dehydrogenase (15-hydroxy PGDH), but it is not a substrate for the enzyme (North et al.; Ohno et al). Because of its resistance to metabolism by 15-hydroxy PGDH, 16,16-Dimethyl PGE2 has a prolonged half-life in vivo. 16,16-Dimethyl PGE2 acts as an agonist on most prostaglandin E (EP) receptor subtypes (Coleman et al.; Robert et al.). The Kd for activation of isolated EP2 receptors is about 1 nM (Coleman et al.). 16,16-Dimethyl PGE2 is supplied in methyl acetate solution at 10 mg/mL (26 mM).

Molecular Name:	16,16-Dimethyl Prostaglandin E2
Alternative Names:	16,16-dimethyl PGE2
CAS Number:	39746-25-3; 79-20-9
Chemical Formula:	C ₂₂ H ₃₆ O ₅
Molecular Weight:	380.5 g/mol
Purity:	≥ 98%
Chemical Name:	9-oxo-11α,15R-dihydroxy-16,16-dimethyl-prosta-5Z,13E-dien-1-oic acid
Structure:	



Properties

Physical Appearance:	A solution in methyl acetate
Storage:	Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com .
Solubility:	Not applicable.

Published Applications

MAINTENANCE AND SELF-RENEWAL

- Increased hematopoietic stem and progenitor cell (HSPC) numbers in zebrafish aorta-gonad-mesonephros (AGM) region and mouse bone marrow (North et al.).
- Mediates the effects of WNT on zebrafish HSPC self-renewal (Goessling et al.).

References

- Coleman RA et al. (1994) International Union of Pharmacology classification of prostanoid receptors: properties, distribution, and structure of the receptors and their subtypes. *Pharmacol Rev* 46(2): 205–29.
- Goessling W et al. (2009) Genetic interaction of PGE2 and Wnt signaling regulates developmental specification of stem cells and regeneration. *Cell* 136(6): 1136–47.
- North TE et al. (2007) Prostaglandin E2 regulates vertebrate haematopoietic stem cell homeostasis. *Nature* 447(7147): 1007–11.
- Ohno H et al. (1978) Studies on 15-Hydroxyprostaglandin Dehydrogenase with Various Prostaglandin Analogues. *J Biochem* 84(6): 1485–1494.
- Robert A et al. (1976) Gastric antisecretory and antiulcer properties of PGE2, 15-methyl PGE2, and 16, 16-dimethyl PGE2. Intravenous, oral and intrajejunal administration. *Gastroenterology* 70(3): 359–70.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.