

## Small Molecules

### Cardiogenol C

Inducer of cardiac differentiation

Catalog # 72422  
72424

1 mg  
10 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## Product Description

Cardiogenol C is a diaminopyrimidine that induces cardiomyogenesis in mouse embryonic stem cells. This product is supplied as the hydrochloride salt of the molecule.

Molecular Name:	Cardiogenol C (Hydrochloride)
Alternative Names:	Cardiogenol C; Cardiogenol C hydrochloride
CAS Number:	671225-39-1
Chemical Formula:	C <sub>13</sub> H <sub>16</sub> N <sub>4</sub> O <sub>2</sub> · HCl
Molecular Weight:	296.8 g/mol
Purity:	≥ 97%
Chemical Name:	2-((2-((4-methoxyphenyl)amino)pyrimidin-4-yl)amino)ethan-1-ol hydrochloride
Structure:	



## Properties

Physical Appearance:	A crystalline solid
Storage:	Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com.
Solubility:	<ul style="list-style-type: none"><li>· PBS (pH 7.2) ≤ 30 mM</li><li>· DMSO ≤ 65 mM</li><li>· Absolute ethanol ≤ 3 mM</li></ul> For example, to prepare a 10 mM stock solution in PBS, resuspend 1 mg in 337 μL of PBS (pH 7.2).

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

## Published Applications

### REPROGRAMMING

- Induces trans-differentiation of mouse CD34+K15+ hair bulge progenitor cells into cardiomyocyte-like cells (Yau et al.).
- Induces cardiomyogenic function in the lineage-committed progenitor cells, C2C12 skeletal myoblasts and mouse A5 cardiovascular progenitor cells (Mike et al.).

### DIFFERENTIATION

- Induces the differentiation of myosin heavy chain (MHC)-positive cardiomyocytes from mouse embryonic stem cells (Wu et al.).

## References

Mike AK et al. (2014) Small molecule cardiogenol C upregulates cardiac markers and induces cardiac functional properties in lineage-committed progenitor cells. *Cell Physiol Biochem* 33(1): 205–21.

Wu X et al. (2004) Small molecules that induce cardiomyogenesis in embryonic stem cells. *J Am Chem Soc* 126(6): 1590–1.

Yau WW et al. (2011) Cardiogenol C can induce Mouse Hair Bulge Progenitor Cells to Transdifferentiate into Cardiomyocyte-like Cells. *Proteome Sci* 9(1): 3.

## Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at [www.stemcell.com/smallmolecules](http://www.stemcell.com/smallmolecules) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2015 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.