IDE1

Small Molecules

Activin/BMP/TGF-β pathway activator

STEMCELLTM TECHNOLOGIES

Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog # 72512 1 mg 72514 5 mg

Product Description

Inducer of Definitive Endoderm 1 (IDE1) induces differentiation of mouse or human embryonic stem (ES) cells by activating SMAD2 phosphorylation and NODAL expression (Borowiak et al.). At $EC_{50} = 125$ nM, SOX17 expression was induced in mouse ES cells.

Molecular Name: IDE1

Alternative Names: Not applicable CAS Number: 1160927-48-9 Chemical Formula: $C_{15}H_{18}N_2O_5$ Molecular Weight: 306.3 g/mol Purity: $\geq 95\%$

Chemical Name: 1-[2-[(2-carboxyphenyl)methylene]hydrazide]-heptanedioic acid

Structure:

$$HO_2C$$
 N
 H
 N
 H
 N
 H

Properties

Physical Appearance: A crystalline solid

Storage: Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please

contact techsupport@stemcell.com.

Solubility: · Absolute ethanol ≤ 320 µM

· DMSO ≤ 80 mM

For example, to prepare a 10 mM stock solution in DMSO, resuspend 1 mg in 326 µL of fresh DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Small Molecules IDE1



Published Applications

DIFFERENTIATION

· Induces differentiation of mouse or human ES cells to definitive endoderm in the absence of Activin A, NODAL, or feeder cells (Borowiak et al.).

References

Borowiak M et al. (2009) Small molecules efficiently direct endodermal differentiation of mouse and human embryonic stem cells. Cell Stem Cell 4(4): 348–58.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2015 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.