
Small Molecules	Gatifloxacin	STENCELL ^M
	Antibiotic; Inhibits bacterial DNA gyrase and topoisomerase IV	Scientists Helping Scientists [™] WWW.STEMCELL.COM
Catalog # 72752	1 g	TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Gatifloxacin is a fluoroquinolone antibiotic which inhibits bacterial DNA gyrase (IC₅₀ = $0.109 \,\mu$ g/ml) and topoisomerase IV (IC₅₀ = 13.8 µg/ml; Takei et al.). It is much less effective against HeLa cell topoisomerase II (IC₅₀ = 265 µg/ml; Takei et al.).

Molecular Name:	Gatifloxacin	
Alternative Names:	AM 1155; BMS 206584-01; PD 135432	
CAS Number:	112811-59-3	
Chemical Formula:	$C_{19}H_{22}FN_3O_4$	
Molecular Weight:	375.4 g/mol	
Purity:	≥ 98%	
Chemical Name:	1-Cyclopropyl-6-fluoro-8-methoxy-7-(3-methylpiperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid	
Structure:	O O	



Properties

Physical Appearance:	A crystalline solid
Storage:	Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com.
Solubility:	· Absolute ethanol \leq 20 mM

Absolute ethanol \leq 20 mM

 \cdot DMSO \leq 45 mM

For example, to prepare a 20 mM stock solution in DMSO, resuspend 1 mg in 225 µL of fresh DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.



Published Applications

MAINTENANCE AND SELF-RENEWAL

· Promotes self-renewal in cultured human and mouse embryonic stem (ES) cells (Desbordes et al.).

References

Desbordes SC et al. (2008) High-throughput screening assay for the identification of compounds regulating self-renewal and differentiation in human embryonic stem cells. Cell Stem Cell 2(6): 602–12. Takei M et al. (1998) Inhibitory Activities of Gatifloxacin (AM-1155), a Newly Developed Fluoroquinolone, against Bacterial and Mammalian Type II Topoisomerases. Antimicrob Agents Chemother 42(10): 2678–2681.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2015 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.