

## Antibodies

### Mouse IgG2b, kappa Isotype Control Antibody, Clone MPC-11, Biotin

Mouse monoclonal IgG2b, kappa isotype control antibody, biotin-conjugated

Catalog #60072BT  
#60072BT.1

200 µg 0.5 mg/mL  
50 µg 0.5 mg/mL



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## Product Description

The MPC-11 antibody (IgG2b, kappa) is suitable for use as an isotype-matched control antibody in several applications to estimate the degree of non-specific binding by an antigen-specific antibody. Ideally, the isotype control should have the same subclass of heavy chain (IgA, IgD, IgE, IgG, or IgM) and light chain (kappa or lambda) as the specific antibody being employed. If a conjugated antibody is employed, an isotype control conjugated to the same molecule (e.g. fluorochrome) should be chosen. The use of an appropriate isotype control helps confirm the specificity of the antigen-specific antibody and indicates non-specific binding that may result from binding to Fc receptors or other cell components. The myeloma cell line secreting the MPC-11 antibody was derived from the transplantable Merwin Plasmacytoma-11 cell line, which was induced by implantation of a plastic diffusion chamber into a BALB/c mouse. This product has unknown binding specificity, having been screened on a variety of activated, resting, live and fixed tissues from several species, including mouse, rat, human, and non-human primates.

Target Antigen Name:	IgG2b Isotype Control
Alternative Names:	Not applicable
Gene ID:	Not applicable
Species Reactivity:	Not applicable
Host Species:	Mouse (BALB/c)
Clonality:	Monoclonal
Clone:	MPC-11
Isotype:	IgG2b, kappa
Immunogen:	Not applicable
Conjugate:	Biotin

## Applications


Verified:	FC
Reported:	FC
Special Applications:	This antibody clone has been verified for use as an isotype control antibody for assessing non-specific binding to cells in flow cytometry and immunofluorescence microscopy applications (surface and intracellular staining).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

## Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Purification:	The antibody was purified by affinity chromatography and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. For product expiry date, please contact techsupport@stemcell.com.
Directions for Use:	The suggested use of this antibody is at concentrations comparable to those of the specific antibody of interest.

## Data



Flow cytometry analysis of human peripheral blood mononuclear cells (PBMCs) labeled with Mouse IgG2b, kappa Isotype Control Antibody, Clone MPC-11, Biotin, followed by streptavidin (SAV) APC (solid line histogram). Filled histogram shows labeling with a mouse IgG2b, kappa positive control antibody (Anti-Human CD20 Antibody, Clone 2H7, Biotin; Catalog #60008BT), followed by SAV APC.

## Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at [www.stemcell.com/antibodies](http://www.stemcell.com/antibodies) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## References

1. Jafari M et al. (2014) Activities of transmitted/founder and chronic clade B HIV-1 Vpu and a C-terminal polymorphism specifically affecting virion release. *J Virol* 88(9): 5062–78. (FC)
2. Le Saout C et al. (2014) Chronic exposure to type-I IFN under lymphopenic conditions alters CD4 T cell homeostasis. *PLoS Pathog* 10(3): e1003976. (FA)
3. Reynolds A. E. et al. (2014) Natural IgM is produced by CD5- plasma cells that occupy a distinct survival niche in bone marrow. *J Immunol* 194(1): 231–42. (FC)
4. Wolpert F et al. (2014) A disintegrin and metalloproteinases 10 and 17 modulate the immunogenicity of glioblastoma-initiating cells. *Neuro Oncol* 16(3): 382–91. (FC)
5. Berger V et al. (2013) An anti-TNFR1 scFv-HSA fusion protein as selective antagonist of TNF action. *Protein Eng Des Sel* 26(10): 581–87. (FC)
6. Zhou H et al. (2013) Fibrinogen-specific antibody induces abdominal aortic aneurysm in mice through complement lectin pathway activation. *Proc Natl Acad Sci USA* 110(46): E4335–44. (FA, IHC)
7. Zhou X et al. (2013) Variation in dietary salt intake induces coordinated dynamics of monocyte subsets and monocyte-platelet aggregates in humans: implications in end organ inflammation. *PLoS One* 8(4): e60332. (FC)
8. Der-Petrossian M et al. (2011) Dermal infiltrates of cutaneous T-cell lymphomas with epidermotropism but not other cutaneous lymphomas are abundant with langerin + dendritic cells. *J Eur Acad Dermatol Venerol* 25(8): 922–27. (IF, IHC)
9. Podolin PL et al. (2008) Inhibition of invariant chain processing, antigen-induced proliferative responses, and the development of collagen-induced arthritis and experimental autoimmune encephalomyelitis by a small molecule cysteine protease inhibitor. *J Immunol* 180(12): 7989–8003. (FC, WB)
10. Spetz A et al. (2008) IgG regulates the CD1 expression profile and lipid antigen-presenting function in human dendritic cells via FcγRIIIa. *Blood* 111(10): 5037–46. (FA/Blocking)
11. Day CL et al. (2006) PD-1 expression on HIV-specific T cells is associated with T-cell exhaustion and disease progression. *Nature* 443(7106): 350–54. (FA/Blocking)
12. Kumamoto T et al. (1999) Human dendritic cells express the thrombopoietin receptor, c-Mpl. *Br J Haematol* 105(4): 1025–33. (FC, ICC)

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Canada Inc. E8 and TeSR are trademarks of WARF. Vitronectin XF is developed and manufactured by Primorigen Biosciences, Inc. and Vitronectin XF is a trademark of Primorigen Biosciences, Inc. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. This product is licensed for internal research use only and its sale is expressly conditioned on the buyer not using it for manufacturing, performing a service, or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or [outlicensing@lifetech.com](mailto:outlicensing@lifetech.com). While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.