

ReproRNA™-OKSGM Kit

Kit for generating iPS cells using ReproRNA™-OKSGM,
a non-integrating, self-replicating RNA reprogramming vector



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

ReproRNA™-OKSGM is a non-integrating, self-replicating RNA-based reprogramming vector for generating induced pluripotent stem (iPS) cells. This vector expresses Oct-3/4, Klf-4, Sox2, Glis1, c-Myc, and a puromycin-resistant cassette in a single construct. The ReproRNA™-OKSGM Kit includes ReproRNA™-OKSGM vector, Recombinant B18R Protein, ReproRNA™ Transfection Reagent, and ReproRNA™ Transfection Supplement. ReproRNA™-OKSGM vector (Catalog #05931) and Recombinant B18R Protein (Catalog #78075) are also available for individual sale.

- ReproRNA™-OKSGM is an RNA-based reprogramming system
- Non-integrating
- Non-viral
- Single transfection

Ordering Information

PRODUCT NAME	CATALOG #	SIZE	KIT COMPONENTS
ReproRNA™-OKSGM Kit	05930	1 Kit	<ul style="list-style-type: none">• ReproRNA™-OKSGM• Recombinant B18R Protein• ReproRNA™ Transfection Reagent• ReproRNA™ Transfection Supplement
ReproRNA™-OKSGM	05931	12 µg (1 µg/µL)	Not applicable.
Recombinant B18R Protein	78075	50 µg	Not applicable.

Component Storage and Stability

COMPONENT NAME	CATALOG #	SIZE	STORAGE	SHELF LIFE
ReproRNA™-OKSGM	05931	12 µg	Store at -80°C.**	Stable for 24 months from date of manufacture (MFG) on label.
Recombinant B18R Protein	78075	50 µg	Store at -80°C.	Stable as supplied for 12 months from date of receipt.
ReproRNA™ Transfection Reagent*†	05932	100 µL	Store at 2 - 8°C.	Stable until expiry date (EXP) on label.
ReproRNA™ Transfection Supplement*†	05933	100 µL	Store at 2 - 8°C.	Stable until expiry date (EXP) on label.

**Do not store at -20°C.

*This component is sold as part of the ReproRNA™-OKSGM Kit (Catalog #05930) and is not available for individual sale.

†Please refer to the Safety Data Sheet (SDS) for hazard information.

Materials Required But Not Included

PRODUCT NAME	CATALOG #
Corning® Matrigel® hESC-Qualified Matrix	Corning 354277
Falcon® 6-Well Flat-Bottom Plate, Tissue Culture-Treated	38016
D-PBS (Without Ca ⁺⁺ and Mg ⁺⁺)	37350
DMEM with 4500 mg/L D-Glucose	36250
Fetal bovine serum (FBS)	---
MEM Non-Essential Amino Acid Solution (100X)	07600
L-Glutamine	07100
Advanced DMEM	Life Technologies 12491-015
Opti-MEM® I Reduced-Serum Medium	Life Technologies 31985-062
ReproTeSR™	05920
Puromycin	73342
Trypsin-EDTA (0.25%)	07901
Sterile microtubes	Sarstedt 72.730.005
iPS cell maintenance medium (e.g. mTeSR™1 or TeSR™-E8™)	85850 or 05940

Preparation of Reagents and Materials

Use sterile techniques to prepare the following reagents and materials.

1) FIBROBLAST CULTURE MEDIUM

The following example is for preparing 500 mL of Fibroblast Culture Medium. If preparing other volumes, adjust accordingly.

Combine the following:

- 440 mL DMEM with 4500 mg/L D-Glucose
- 50 mL FBS
- 5 mL MEM Non-Essential Amino Acid Solution (100X)
- 5 mL 200 mM L-Glutamine

Pre-warm to room temperature (15 - 25°C) before use. Store Fibroblast Culture Medium at 2 - 8°C for up to 2 weeks.

2) GROWTH MEDIUM

The following example is for preparing 70 mL of Growth Medium (sufficient for 6 wells). If preparing other volumes, adjust accordingly.

Combine the following:

- 62.3 mL Advanced DMEM
- 7 mL FBS
- 700 µL 200 mM L-Glutamine
- 24.5 µL 0.5 mg/mL Recombinant B18R Protein (final concentration 175 ng/mL)

Warm to room temperature (15 - 25°C) before use. Store Growth Medium at 2 - 8°C for up to 1 week.

3) ReproRNA™ COCKTAIL

NOTE: Prepare ReproRNA™ Cocktail immediately before transfection.

ReproRNA™-OKSGM is an RNA-based vector; use RNase- and DNase-free pipette tips and microtubes when preparing the ReproRNA™ Cocktail.

For each well (6-well plate format) of fibroblasts to be reprogrammed, combine the following components in the order shown in a sterile microtube:

- 1 µL ReproRNA™-OKSGM
- 100 µL Opti-MEM® 1 Reduced-Serum Medium
- 2 µL ReproRNA™ Transfection Supplement
- 2 µL ReproRNA™ Transfection Reagent

Pipette gently to mix after adding each component. Incubate ReproRNA™ Cocktail at room temperature (15 - 25°C) for exactly 5 minutes before immediately adding to cells. Volumes can be scaled up when reprogramming multiple wells of fibroblasts.

4) GROWTH MEDIUM + PUROMYCIN

NOTE: Prepare Growth Medium + Puromycin fresh daily, as required.

The following example is for preparing 9 mL of Growth Medium + Puromycin (sufficient for 6 wells). If preparing other volumes, adjust accordingly.

Combine the following:

- 7.2 µL of a 1 mg/mL solution of Puromycin (final concentration 0.8 µg/mL)
- 9 mL Growth Medium

5) ReproTeSR™ (with and without B18R)

NOTE: For preparation of complete ReproTeSR™ medium (without B18R), refer to the ReproTeSR™ Product Information Sheet (Document #DX20217), available at www.stemcell.com or contact us to request a copy.

The following example is for preparing 70 mL of ReproTeSR™ with B18R (sufficient for 6 wells). If preparing other volumes, adjust accordingly.

Combine the following:

- 24.5 µL 0.5 mg/mL Recombinant B18R Protein (final concentration 175 ng/mL)
- 70 mL complete ReproTeSR™ medium

Warm to room temperature (15 - 25°C) before use. Store ReproTeSR™ with B18R at 2 - 8°C for up to 1 week.

Protocol Diagram




Figure 1. Timeline for Reprogramming with ReproRNA™-OKSGM

Somatic cells are transfected with ReproRNA™-OKSGM at day 0, and cultured in Growth Medium (containing Puromycin). After 5 days of Puromycin selection post-transfection, cells are cultured in ReproTeSR™ for the remainder of the reprogramming induction phase until iPS cell colonies emerge. Recombinant B18R Protein is also added during the first 2 weeks after transfection to inhibit the interferon response and increase cell viability. Typically, by day 20, iPS cell colonies are large enough to be isolated and propagated in TeSR™ media*.

*TeSR™ family media (mTeSR™1, TeSR™2, or TeSR™-E8™)

Directions for Use

Please read the entire protocol before proceeding. Use sterile techniques when performing the following protocols.

A. CULTURE OF HUMAN DERMAL FIBROBLASTS

Use low-passage (passage 2 - 5) human dermal fibroblasts for reprogramming experiments. Extended passaging of fibroblasts will decrease reprogramming efficiency.

The following instructions are for 1 well of a 6-well plate. If using other cultureware, adjust volumes accordingly.

1. Plate low-passage fibroblasts (3 - 5000 cells/cm²) in Fibroblast Culture Medium and incubate at 37°C and 5% CO₂.
2. Every 2 days, aspirate medium and replace with 2 mL fresh Fibroblast Culture Medium.
3. When the culture is approximately 85% confluent, passage with Trypsin-EDTA (0.25%) as follows:
 - a. Aspirate Fibroblast Culture Medium. Rinse twice with D-PBS (Without Ca⁺⁺ and Mg⁺⁺).
 - b. Add 1 mL Trypsin-EDTA (0.25%).
 - c. Incubate at 37°C and 5% CO₂ for 2 - 5 minutes or until fibroblasts have detached.
 - d. Add 1 mL Fibroblast Culture Medium to inactivate trypsin. Transfer cell suspension to a conical tube.
 - e. Centrifuge at 300 x *g* for 5 minutes. Remove and discard supernatant and resuspend cells in fresh Fibroblast Culture Medium.
 - f. Plate fibroblasts at a split ratio of 1:4 to 1:6. Incubate at 37°C and 5% CO₂ until ready to reprogram the cells.

B. RNA TRANSFECTION PROTOCOL

The protocol below describes the reprogramming of human dermal fibroblasts in 1 well of a 6-well plate. If using other cultureware, adjust volumes accordingly.

Harvesting and Plating Fibroblasts

1. Coat a tissue culture-treated 6-well plate with Corning® Matrigel® hESC-Qualified Matrix and bring to room temperature (15 - 25°C) for at least 30 minutes prior to use.
NOTE: For complete instructions on coating plates with Corning® Matrigel®, refer to the Technical Manual: Maintenance of Human Pluripotent Stem Cells in mTeSR™1 (Document #28315) or TeSR™-E8™ (Document #29267), available at www.stemcell.com or contact us to request a copy.
2. Harvest fibroblasts (from section A) with Trypsin-EDTA (0.25%) as described in section A, steps 3a - 3d.
3. Centrifuge at 300 x *g* for 5 minutes. Remove and discard supernatant and resuspend cells in fresh Fibroblast Culture Medium at 5 x 10⁴ cells/mL.
4. Add 2 mL of the cell suspension (1 x 10⁵ cells) to 1 well of a pre-coated 6-well plate (prepared in step 1).
5. Incubate at 37°C and 5% CO₂ overnight to allow fibroblasts to attach to the matrix.

Transfection

DAY 0

6. Aspirate Fibroblast Culture Medium from each well and wash with 2 mL D-PBS (Without Ca⁺⁺ and Mg⁺⁺) per well.
7. Add 1 mL Growth Medium to each well. Incubate at 37°C and 5% CO₂ for 20 minutes while preparing ReproRNA™ Cocktail (step 8).
8. Prepare ReproRNA™ Cocktail in a sterile microtube for each well of fibroblasts to be reprogrammed. Incubate at room temperature (15 - 25°C) for 5 minutes.
9. Immediately add ReproRNA™ Cocktail dropwise to each well containing fibroblasts. Gently rock the plate using a back-and-forth and side-to-side motion to ensure the ReproRNA™ Cocktail is evenly distributed throughout the entire well.
10. Incubate at 37°C and 5% CO₂ overnight.

DAY 1

11. Aspirate Growth Medium containing ReproRNA™ Cocktail and add 1.5 mL Growth Medium + Puromycin per well.
NOTE: The addition of Growth Medium + Puromycin removes fibroblasts that have not been transfected with ReproRNA™-OKSGM.
12. Incubate at 37°C and 5% CO₂ overnight.

DAY 2 - 5

13. Perform a daily medium change with 1.5 mL of fresh Growth Medium + Puromycin per well. Monitor fibroblasts for cell morphology and survival. Incubate at 37°C and 5% CO₂.

DAY 6 - 7

14. Perform a daily medium change with 1.5 mL Growth Medium (without Puromycin) per well. Incubate at 37°C and 5% CO₂.

DAY 8

15. Aspirate Growth Medium and add 1.5 mL ReproTeSR™ **with** B18R per well. Incubate at 37°C and 5% CO₂ overnight.

DAY 9 - 14

16. Perform a daily medium change with ReproTeSR™ **with** B18R. Incubate at 37°C and 5% CO₂.

DAY 15 - 28

17. Perform a daily medium change with ReproTeSR™ (**without** B18R) until iPS colonies form and are ready to be manually isolated. iPS colonies typically arise between 15 - 28 days post-transfection of ReproRNA™-OKSGM.

18. Manually isolate putative iPS cell colonies. Use either a 22 - 25 gauge needle or a pulled glass pipette to cut the putative iPS cell colony into small fragments. Then use a 200 µL pipettor with a filtered tip to scrape and aspirate colony fragments.

NOTE: If there are many untransfected, partially reprogrammed and/or differentiated cells surrounding the putative iPS cell colony, these may need to be scraped away prior to isolating the iPS cell colony.

19. Immediately plate iPS cell colony fragments on cultureware coated with desired matrix (e.g. Corning® Matrigel®) and containing iPS cell maintenance medium (e.g. mTeSR™1 or TeSR™-E8™).

NOTE: To facilitate the initial attachment of iPS cell colony fragments, add Y-27632 (Catalog #72302) to the maintenance medium at a final concentration of 10 µM. After 24 hours, replace the maintenance medium (without Y-27632).

20. Incubate at 37°C and perform iPS cell maintenance medium changes as appropriate.

NOTE: For complete instructions on how to maintain iPS cells using mTeSR™1 or TeSR™-E8™, refer to the Technical Manual: Maintenance of Human Pluripotent Stem Cells in mTeSR™1 (Document #28315) or TeSR™-E8™ (Document #29267), available at www.stemcell.com or contact us to request a copy.

Assessment of hPSCs

The following antibodies can be used to characterize hPSCs by flow cytometry or immunocytochemistry:

- Anti-Human SSEA-4 Antibody, Clone MC-813-70 (Catalog #60062)
- Anti-Human TRA-1-60 Antibody, Clone TRA-1-60R (Catalog #60064)
- Anti-Human OCT4 (OCT3) Antibody, Clone 3A2A20 (Catalog #60093)

Related Products

For related products, including specialized cell culture and storage media, matrices, antibodies, cytokines, and small molecules, visit www.stemcell.com/hPSCworkflow or contact us at techsupport@stemcell.com.

References

Yoshioka N et al. (2013) Efficient generation of human iPSCs by a synthetic self-replicative RNA. *Cell Stem Cell* 13(2): 246–54.

(a) This Product/Service shall be used for customers' internal research; (b) This Product/Service shall not be used or administered in (1) human subjects for human clinical use or (2) animals for veterinary use for therapeutic, diagnostic or prophylactic purposes, including but not limited to clinical applications, cell therapy, transplantation, and/or regenerative medicines; (c) If a customer wishes to conduct the following commercial use and he/she has not taken any appropriate patent license from AJ, such customer needs to contact iPS Academia Japan, Inc. ("AJ") and take a commercial license from AJ to cover such commercial use: (1) use of Product/Service and/or iPS cells generated by using Product for manufacture of related products for sale, including but not limited to culture medium and equipment. (2) use of Product/Service, iPS cells generated and/or their modifications generated by using Product/Service for screening compounds, antibodies, proteins, peptides for commercial use, except for target discovery, target validation or assay development. (3) sale, lease, distribution, transfer of iPS cells and/or modifications generated by using Product/Service or use such to provide a service, information or data, to a third party for financial gain; (d) In case that a customer sale, lease, distribution, transfer of iPS cells generated by using Product/Service to for-profit organization not for financial gain, such customer shall notify the for-profit organization who receives the iPS cells generated by using Product/Service that such for-profit organization is required to take a license from AJ to use them for internal research of any kind; (e) In any case that a customer transfers the Product/Service, iPS cells and/or modifications generated by using such Product/Service, the customer shall notify the receipt regarding this Limited Use Label License and provide a copy of it together with such material transfer; (f) Other than rights granted herein, no other right, express or implied, is conveyed by the sale of Product or the iPS cells generated by using Product/Service or the provision of generation service of such iPS cells.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, ReproTeSR, and ReproRNA are trademarks of STEMCELL Technologies Canada Inc. Corning, Falcon, and Matrigel are registered trademarks of Corning Incorporated. Opti-MEM is a registered trademark of Thermo Fisher Scientific. mTeSR, TeSR, and E8 are trademarks of WARF. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.