

## Anti-Mouse CD45.2 Antibody, Clone 104, Biotin



Scientists Helping Scientists™ | [WWW.STEMCELL.COM](http://WWW.STEMCELL.COM)

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

[INFO@STEMCELL.COM](mailto:INFO@STEMCELL.COM) • [TECHSUPPORT@STEMCELL.COM](mailto:TECHSUPPORT@STEMCELL.COM)

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## Antibodies

Mouse monoclonal IgG2a antibody  
against mouse CD45.2, biotin-  
conjugated

Catalog #60118BT  
#60118BT.1

500 µg 0.5 mg/mL  
50 µg 0.5 mg/mL

## Product Description

The 104 monoclonal antibody reacts with mouse alloantigen CD45.2 (Ly5.2) expressed by all hematopoietic cells except mature erythrocytes and platelets in mouse strains A, AKR, BALB/c, CBA/Ca, CBA/J, C3H/He, C57BL, C57BR, C57L, C58, DBA/1, DBA/2, NZB, SWR, and 129. Clone 104 does not react with leukocytes that express the CD45.1 alloantigen. CD45.2 is an allelic form of CD45 (Leukocyte Common Antigen), a 180 - 240 kD type I transmembrane glycoprotein. CD45 is a member of the protein tyrosine phosphatase (PTP) family and contains two catalytic domains in the intracellular (COOH) domain, which regulate cellular signaling. The extracellular domains are highly variable, due to alternative splicing among exons 4, 5, and 6, and differential glycosylation. These CD45 isoforms correlate to specific cell types, as well specific activation and maturation states of immune cells. The 104 monoclonal antibody has been used extensively in adoptive cell transfer and bone marrow transplantation studies in mice, allowing for the differentiation of CD45.2- and CD45.1-expressing cells.

Target Antigen Name:	CD45.2
Alternative Names:	B220, CD45 antigen, CD45R, GP180, LCA, Leukocyte common antigen, Ly5.2, Protein tyrosine phosphatase receptor type C, PTPRC, T200
Gene ID:	19264
Species Reactivity:	Mouse, does not react with mouse CD45.1
Host Species:	Mouse (SJL)
Clonality:	Monoclonal
Clone:	104
Isotype:	IgG2a, kappa
Immunogen:	B10.S mouse thymocytes and splenocytes
Conjugate:	Biotin

## Applications


Verified:	FC
Reported:	FC

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

## Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Purification:	The antibody was purified by affinity chromatography and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. For product expiry date, please contact <a href="mailto:techsupport@stemcell.com">techsupport@stemcell.com</a> .
Directions for Use:	For flow cytometry the suggested use of this antibody is $\leq 0.25 \mu\text{g}$ per $1 \times 10^6$ cells in 100 µL. It is recommended that the antibody be titrated for optimal performance for each application.

## Data



Flow cytometry analysis of C57BL/6 mouse splenocytes labeled with Anti-Mouse CD45.2 Antibody, Clone 104, Biotin, followed by streptavidin (SAV) APC (filled histogram) or Mouse IgG2a, kappa Isotype Control Antibody, Clone MOPC-173, Biotin (Catalog #60071BT), followed by SAV APC (solid line histogram).

## Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at [www.stemcell.com/antibodies](http://www.stemcell.com/antibodies) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## References

1. Mackay LK et al. (2015) Cutting edge: CD69 interference with sphingosine-1-phosphate receptor function regulates peripheral T cell retention. *J Immunol* 194(5): 2059–63. (FC)
2. Rabenhorst U et al. (2015) Single-stranded DNA-binding transcriptional regulator FUBP1 is essential for fetal and adult hematopoietic stem cell self-renewal. *Cell Rep* 11(12): 1847–55. (FACS, FC)
3. Tait Wojno ED et al. (2015) The prostaglandin D2 receptor CRTH2 regulates accumulation of group 2 innate lymphoid cells in the inflamed lung. *Mucosal Immunol* 8(6): 1313–23. (FC)
4. van Blijswijk J et al. (2015) Altered lymph node composition in diphtheria toxin receptor-based mouse models to ablate dendritic cells. *J Immunol* 194(1): 307–15. (FC)
5. Nakano-Yokomizo T et al. (2011) The immunoreceptor adapter protein DAP12 suppresses B lymphocyte-driven adaptive immune responses. *J Exp Med* 208(8): 1661–71. (FC)
6. Gebhardt T et al. (2009) Memory T cells in nonlymphoid tissue that provide enhanced local immunity during infection with herpes simplex virus. *Nat Immunol* 10(5): 524–30. (FC)
7. Trowbridge JJ et al. (2009) DNA methyltransferase 1 is essential for and uniquely regulates hematopoietic stem and progenitor cells. *Cell Stem Cell* 5(4): 442–9. (FC)
8. Kohlmeier JE et al. (2008) The chemokine receptor CCR5 plays a key role in the early memory CD8+ T cell response to respiratory virus infections. *Immunity* 29(1): 101–13. (FC)
9. Phan TG et al. (2007) Subcapsular encounter and complement-dependent transport of immune complexes by lymph node B cells. *Nat Immunol* 8(9): 992–1000. (FC, IF)
10. Duncan AW et al. (2005) Integration of Notch and Wnt signaling in hematopoietic stem cell maintenance. *Nat Immunol* 6(3): 314–22. (FC)
11. Hock H et al. (2004) Gfi-1 restricts proliferation and preserves functional integrity of haematopoietic stem cells. *Nature* 431(7011): 1002–7. (FC)
12. Wilson A et al. (2001) Notch 1-deficient common lymphoid precursors adopt a B cell fate in the thymus. *J Exp Med* 194(7): 1003–12. (FC)
13. Shen FW et al. (1986) Further definition of the Ly-5 system. *Immunogenetics* 24(3): 146–9.
14. Yakura H et al. (1983) On the function of Ly-5 in the regulation of antigen-driven B cell differentiation. Comparison and contrast with Lyb-2. *J Exp Med* 157(4): 1077–88.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. Antibodies conjugated to Alexa Fluor® are licensed for internal research use only and sale is expressly conditioned on the buyer not using the antibody for manufacturing, performing a service or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or [outlicensing@lifetech.com](mailto:outlicensing@lifetech.com). While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.