

Cytokines

Mouse Recombinant IFN-gamma

Interferon-gamma

Catalog #	78021.1	20 µg
	78021	100 µg
	78021.2	1000 µg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Interferon-gamma (IFN- γ), also known as type II interferon, is produced by T and NK cells, and in smaller amounts by dendritic cells and macrophages. IFN- γ is controlled by cytokines such as IL-12 and IL-18 that are secreted in response to infection (Schroder et al.). IFN- γ binds to a receptor complex and initiates signal transduction via the JAK/STAT pathway; this culminates in the transcription and activation of many genes that control a diverse array of immunological functions (de Weerd & Nguyen; Krause et al.). IFN- γ stimulates the antimicrobial and anti-tumor activity of macrophages, NK cells, and neutrophils (Billiau & Matthys) by promoting the activation of microbial effector functions such as production of reactive oxygen species, NO intermediates, complement, etc. (Schroder et al.). IFN- γ enhances MHC class I and II expression in dendritic cells and mononuclear phagocytes, as well as the production of IL-12 by dendritic cells. In B cells, IFN- γ stimulates survival and growth in both mouse and human cells, and redirects B cells from proliferation towards differentiation. IFN- γ favors the development of Th1 vs Th2 cells and stimulates monocyte differentiation and function (Schroder et al.).

Product Information

Alternative Names:	Interferon gamma, Type II interferon
Accession Number:	P01580
Amino Acid Sequence:	MHGTVIESLE SLNNYFNSSG IDVEEKSLFL DIWRNWQKDG DMKILQSQII SFYLRLEFVL KDNQAISNNI SVIESHLITT FFSNSKAKKD AFMSIAKFEV NNPQVQRQAF NELIRVVHQL LPESSLRKRK RSRC
Predicted Molecular Mass:	15.7 kDa
Species:	Mouse
Cross Reactivity:	Human, Rat
Formulation:	Lyophilized from a sterile-filtered aqueous solution containing sodium phosphate and sodium chloride, pH 7.5.
Source:	E. coli

Specifications


Activity:	The specific activity is $\geq 1 \times 10^7$ units/mg after normalizing to an internal standard as determined in a viral challenge assay using encephalomyocarditis (EMC) virus on L929 cells.
Purity:	$\geq 95\%$
Endotoxin Level:	Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is ≤ 1 EU/ μ g protein.

Preparation and Storage

Storage:	Store at -20°C to -80°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Bring vial and sterile water to room temperature (15 - 25°C). Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Let solution sit for 1 minute at room temperature (15 - 25°C). If precipitate is observed, centrifuge at 16,000 x g for 1 minute. Remove supernatant and transfer to a new tube, taking care not to disturb the pellet. Discard the pellet. A 10% overfill has been added to compensate for any loss of protein in the precipitate.

OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide, do not store at 2 - 8°C for more than 1 month or at -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Mouse Recombinant IFN-gamma was measured with L929 cells exposed to the EMC virus in a cytopathic effect (CPE) assay. Cell survival was measured using a fluorometric assay method. The EC₅₀ is defined as the effective concentration of the cytokine at which cell survival is at 50% of maximum. The EC₅₀ in the above graph is 0.22 ng/mL. When normalized to an internal standard, the specific activity in the above example is 4.3×10^7 units/mg.

(B) 1 μ g of Mouse Recombinant IFN-gamma was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Mouse Recombinant IFN-gamma has a predicted molecular mass of 15.7 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Billiau A & Matthys P. (2009) Interferon-gamma: a historical perspective. *Cytokine Growth Factor Rev* 20(2): 97–113.
- Krause CD et al. (2000) Signaling by covalent heterodimers of interferon-gamma. Evidence for one-sided signaling in the active tetrameric receptor complex. *J Biol Chem* 275(30): 22995–3004.
- Schroder K et al. (2004) Interferon-gamma: an overview of signals, mechanisms and functions. *J Leukoc Biol* 75(2): 163–89.
- de Weerd NA & Nguyen T. (2012) The interferons and their receptors--distribution and regulation. *Immunol Cell Biol* 90(5): 483–91.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.