Mouse Recombinant IL-2

Cytokines

Interleukin 2

Catalog # 78081 20 µg

> 78081.1 100 µg 78081.2 1000 µg



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WERSITE

Product Description

Interleukin 2 (IL-2) is a monomeric cytokine that was originally identified as a T cell growth factor (Gaffen & Liu). It binds to a heterotrimeric receptor consisting of CD25, CD122, and CD132. Upon binding, it activates JAK3-, STAT5-, and AKT-dependent signaling pathways, which results in cellular proliferation and survival (Ma et al.). The majority of IL-2 is secreted by activated CD4+ and CD8+ T cells, although B cells and dendritic cells were found to produce IL-2 in small amounts. IL-2 downregulates immune responses to prevent autoimmunity during thymic development, influences the development of CD4+CD25+ regulatory T cells, and affects development of follicular helper T cells. IL-2 also controls inflammation by inhibiting Th17 differentiation (Banchereau et al.). Targeted deletions of the IL-2 gene in mice resulted in development of autoimmune hemolytic anemia, followed by ulcerative colitis. Similar effects were observed in mice that were deficient in IL-2 receptor α (Gaffen & Liu).

Product Information

Alternative Names: Aldesleukin, Interleukin-2, T cell growth factor, TCGF

Accession Number: P04351

Amino Acid Sequence: MAPTSSSTSS STAEAQQQQQ QQQQQQHLE QLLMDLQELL SRMENYRNLK LPRMLTFKFY LPKQATELKD

LQCLEDELGP LRHVLDLTQS KSFQLEDAEN FISNIRVTVV KLKGSDNTFE CQFDDESATV VDFLRRWIAF

CQSIISTSPQ

Predicted Molecular Mass: 17.4 kDa Species: Mouse Cross Reactivity: Human

Formulation: Lyophilized from a sterile-filtered aqueous solution containing sodium citrate, pH 4.0.

Source: E. coli

Specifications

Activity: The specific activity is $\ge 2 \times 10^5$ units/mg (EC50 ≤ 5 ng/mL) as determined by a cell proliferation assay

using CTLL-2 cells.

Purity: ≥ 95%

Endotoxin Level: Measured by kinetic Limulus amebocyte lysate (LAL) analysis and is ≤ 1 EU/µg protein.

Preparation and Storage

Storage: Store at -20°C to -80°C.

Stability: Stable as supplied for 12 months from date of receipt.

Preparation: Centrifuge vial before opening. Reconstitute the product in sterile 10 mM acetic acid to at least 0.1 mg/mL by


pipetting the solution down the sides of the vial. Do not vortex.

OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide, do not store at 2 - 8°C for more than

1 month or at -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.



Data



(A) The biological activity of Mouse Recombinant IL-2 was tested by its ability to promote the proliferation of CTLL-2 cells. Cell proliferation was measured after 3 days of culture using a fluorometric assay method. The EC50 is defined as the effective concentration of the protein at which cell proliferation is at 50% of maximum. The EC50 in the above example is 1.12 ng/mL.

(B) 1 µg of Mouse Recombinant IL-2 was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Mouse Recombinant IL-2 has a predicted molecular mass of 17.4 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

Banchereau J et al. (2012) From IL-2 to IL-37: the expanding spectrum of anti-inflammatory cytokines. Nat Immunol 13(10): 925–31. Gaffen SL & Liu KD. (2004) Overview of interleukin-2 function, production and clinical applications. Cytokine 28(3): 109–23. Ma A et al. (2006) Diverse functions of IL-2, IL-15, and IL-7 in lymphoid homeostasis. Annu Rev Immunol 24: 657–79.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.