

## Anti-Mouse CD27 Antibody, Clone LG.3A10

## Antibodies

Hamster (Armenian) monoclonal IgG1 antibody against human, mouse, rat CD27, unconjugated

Catalog #60160

500 µg 0.5 mg/mL



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## Product Description

The LG.3A10 antibody reacts with CD27, a 45 - 55 kDa member of the tumor necrosis factor receptor superfamily. CD27 is found on peripheral T cells, memory B cells, and NK cells and in mice has been found on most thymocytes, except CD46-CD8- precursors. CD27 and its ligand CD70 are involved in multiple aspects of T cell biology, including the costimulation and expansion of CD4+ and CD8+ T cells and differentiation to memory CD8+ T cells. CD27 interacts with TRAF family members, resulting in enhanced cellular proliferation and the induction of anti-apoptotic proteins. Additionally, CD27 plays important roles in T cell – B cell interactions and the maturation of effector T cells. The expansion of activated B cells and their differentiation to plasma cells is also reported to be enhanced by CD27.

Target Antigen Name:	CD27
Alternative Names:	LPFS2, S152, T14, TNFRSF7, Tp55
Gene ID:	21940/939/500318
Species Reactivity:	Human, Mouse, Rat
Host Species:	Hamster (Armenian)
Clonality:	Monoclonal
Clone:	LG.3A10
Isotype:	IgG1, kappa
Immunogen:	Armenian hamster fibroblast line ARHO12 transfected with mouse Cd27 cDNA
Conjugate:	Unconjugated

## Applications


Verified:	FC
Reported:	FA, FC, IHC, IP
Special Applications:	This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including EasySep™ Human Naive B Cell Isolation Kit (Catalog #17254) and EasySep™ Human Naive B Cell Enrichment Kit (Catalog #19254).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

## Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Purification:	The antibody was purified by affinity chromatography.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. For product expiry date, please contact techsupport@stemcell.com.
Directions for Use:	For flow cytometry the suggested use of this antibody is ≤ 0.25 µg per 1 × 10 <sup>6</sup> cells in 100 µL volume. It is recommended that the antibody be titrated for optimal performance for each application.

## Data



Flow cytometry analysis of C57BL/6 mouse splenocytes labeled with Anti-Mouse CD27 Antibody, Clone LG.3A10, followed by a goat anti-hamster (Armenian) IgG antibody, Alexa Fluor® 488 (filled histogram), or an Armenian hamster IgG isotype control antibody, followed by a goat anti-hamster (Armenian) IgG antibody, Alexa Fluor® 488 (solid line histogram).

## Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at [www.stemcell.com/antibodies](http://www.stemcell.com/antibodies) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## References

1. Kim JH et al. (2016) Programmed cell death ligand 1 alleviates psoriatic inflammation by suppressing IL-17A production from programmed cell death 1-high T cells. *J Allergy Clin Immunol* 137(5): 1466–76.e3. (FC)
2. Chognard G et al. (2014) The dichotomous pattern of IL-12R and IL-23R expression elucidates the role of IL-12 and IL-23 in inflammation. *PLoS One* 9(2): e89092. (FC)
3. Stubelius A et al. (2014) Immunomodulation by the estrogen metabolite 2-methoxyestradiol. *Clin Immunol* 153(1): 40–48. (FC)
4. Vasconcelos JR et al. (2014) Adenovirus vector-induced CD8+ T effector memory cell differentiation and recirculation, but not proliferation, are important for protective immunity against experimental *Trypanosoma cruzi* Infection. *Hum Gene Ther* 25(4): 350–63. (FC)
5. Brodin P et al. (2009) The strength of inhibitory input during education quantitatively tunes the functional responsiveness of individual natural killer cells. *Blood* 113(11): 2434–41. (FC)
6. Hayakawa Y & Smyth MJ. (2006) CD27 dissects mature NK cells into two subsets with distinct responsiveness and migratory capacity. *J Immunol* 176(3): 1517–24. (FC)
7. Arber C et al. (2003) Common lymphoid progenitors rapidly engraft and protect against lethal murine cytomegalovirus infection after hematopoietic stem cell transplantation. *Blood* 102(2): 421–8.
8. Carleton M et al. (2002) Early growth response transcription factors are required for development of CD4(-)CD8(-) thymocytes to the CD4(+)CD8(+) stage. *J Immunol* 168(4): 1649–58. (FC)
9. Gravestain L a et al. (1998) The TNF receptor family member CD27 signals to Jun N-terminal kinase via Traf-2. *Eur J Immunol* 28(7): 2208–16. (FA)
10. Gravestain LA et al. (1996) CD27 cooperates with the pre-T cell receptor in the regulation of murine T cell development. *J Exp Med* 184(2): 675–85. (IHC)
11. Gravestain LA et al. (1995) Novel mAbs reveal potent co-stimulatory activity of murine CD27. *Int Immunol* 7(4): 551–7. (IP)

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.