

## Dissociation Reagents

### Collagenase Type I

For digestion of native collagen fibrils

Catalog # 07415  
07416

100 mg  
1 g



Scientists Helping Scientists™ | [WWW.STEMCELL.COM](http://WWW.STEMCELL.COM)

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

[INFO@STEMCELL.COM](mailto:INFO@STEMCELL.COM) • [TECHSUPPORT@STEMCELL.COM](mailto:TECHSUPPORT@STEMCELL.COM)

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## Product Description

Collagenase is a protease consisting of a single polypeptide chain approximately 1,000 amino acid residues in length. Collagenase is capable of digesting native collagen fibrils commonly found in connective tissues and therefore is frequently used for tissue dissociation. Collagenase Type I contains the activity of several proteases, including collagenase, caseinase, clostripain, and trypsin. Collagenase Type I has been used for the digestion of human tissues such as intestine (Barthel et al.), mammary glands (Huss & Kratz), and prostate (Le et al.), as well as specific cell types such as endothelial cells (Ganguly et al.) and dorsal root ganglion cells (Dib-Hajj et al.).

## Product Information

Alternative Names:	Clostridiopeptidase A; Clostridium histolyticum collagenase; Collagenase 1; Collagenase Type 1; Collagenase I
Format:	Lyophilized powder
Storage:	Store at 2 - 8°C.
Stability:	Stable until expiry date (EXP) on label.
Reconstitution:	Dissociation reagents can be reconstituted in a balanced salt solution or buffer of choice.
Molecular Weight:	68 - 130 kDa
CAS Number:	9001-12-1
Optimum pH:	6.3 - 8.5
Cleavage Site:	-Pro-X-†-Gly-Pro-Y- : X = neutral Y = nonspecific

## Specifications

Source:	Clostridium histolyticum
Activity:	Collagenase: ≥ 125 CDU/mg dry weight (mgdw); Caseinase: ≥ 200 u/mgdw; Clostripain: ≤ 4.0 u/mgdw; Trypsin: ≤ 0.5 u/mgdw. See Notes for further information.

## Related Products

For a complete list of dissociation reagents, as well as related products available from STEMCELL Technologies, please visit our website at [www.stemcell.com](http://www.stemcell.com) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## Notes

### ACTIVITY UNITS

Collagenase: 1 collagenase digestion unit (CDU) equals 1  $\mu\text{mol}$  of L-leucine equivalents released from collagen in 5 hours at 37°C, pH 7.5.

Caseinase: 1 unit equals 1  $\mu\text{mol}$  of L-leucine equivalents released from 25 mg vitamin-free casein in 5 hours at 37°C, pH 7.5. Measures non-specific proteolytic activity.

Clostripain: 1 unit hydrolyzes 1  $\mu\text{mol}$  of N $\alpha$ -benzoyl-L-arginine ethyl ester (BAEE)/minute at 25°C at pH 7.6, after activation in 2.5 mM dithiothreitol (DTT).

Trypsin: 1 unit hydrolyzes 1  $\mu\text{mol}$  of BAEE/minute at 25°C at pH 7.6.

## References

- Barthel ER et al. (2012) Tissue engineering of the intestine in a murine model. *J Vis Exp* (70): e4279.
- Dib-Hajj SD et al. (1999) Two tetrodotoxin-resistant sodium channels in human dorsal root ganglion neurons. *FEBS Lett* 462(1-2): 117–20.
- Ganguly A et al. (2012) Isolation of human umbilical vein endothelial cells and their use in the study of neutrophil transmigration under flow conditions. *J Vis Exp* (66): e4032.
- Huss FRM & Kratz G. (2001) Mammary epithelial cell and adipocyte co-culture in a 3-D matrix: The first step towards tissue-engineered human breast tissue. *Cells Tissues Organs* 169(4): 361–7.
- Le H et al. (2006) DHT and testosterone, but not DHEA or E2, differentially modulate IGF-I, IGFBP-2, and IGFBP-3 in human prostatic stromal cells. *Am J Physiol Endocrinol Metab* 290(5): E952–60.
- Mohapatra A et al. (2016) Group 2 innate lymphoid cells utilize the IRF4-IL-9 module to coordinate epithelial cell maintenance of lung homeostasis. *Mucosal Immunol* 9(1): 275–86.
- Totsuka T et al. (2003) Therapeutic effect of anti-OX40L and anti-TNF-alpha MAbs in a murine model of chronic colitis. *Am J Physiol Gastrointest Liver Physiol* 284(4): G595–603.
- Wang J et al. (2015) Interleukin 18 function in atherosclerosis is mediated by the interleukin 18 receptor and the Na-Cl co-transporter. *Nat Med* 21(7): 820–6.
- Wu Z et al. (2015) Depletion of MEIS2 inhibits osteogenic differentiation potential of human dental stem cells. *Int J Clin Exp Med* 8(5): 7220–30.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.