

Dissociation Reagents

Collagenase A, ACF

Animal component-free collagenase for the digestion of native collagen fibrils

Catalog # 07434
07435

100 mg
1 g



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Collagenase A, Animal Component-Free (ACF) is obtained from cultures free of animal-derived materials. Collagenase is a protease consisting of a single polypeptide chain approximately 1,000 amino acid residues in length. Collagenase is capable of digesting native collagen fibrils commonly found in connective tissues and therefore is frequently used for tissue dissociation. Collagenase preparations contain the activity of several proteases, including collagenase, caseinase, clostripain, and trypsin (Kessler & Yaron). Collagenase A, ACF contains levels of proteolytic activity similar to type I and type II collagenases. Collagenase A, ACF is intended for use in applications where it is necessary to prevent the introduction of potential animal-derived pathogens.

Product Information

Alternative Names:	Clostridium histolyticum collagenase; Collagenase A
Format:	Lyophilized powder
Storage:	Store at 2 - 8°C.
Stability:	Stable until expiry date (EXP) on label.
Reconstitution:	Dissociation reagents can be reconstituted in a balanced salt solution or buffer of choice.
Molecular Weight:	68 - 130 kDa
CAS Number:	9001-12-1
Optimum pH:	6.3 - 8.5
Cleavage Site:	-Pro-X-†-Gly-Pro-Y- : X = neutral Y = nonspecific

Specifications

Source:	Clostridium histolyticum
Activity:	Collagenase: ≥ 150 CDU/mg dry weight (mgdw); Caseinase: ≥ 150 u/mgdw; Clostripain: ≤ 8.0 u/mgdw; Trypsin: ≥ 0.1 u/mgdw. See Notes for further information.

Related Products

For a complete list of dissociation reagents, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com or contact us at techsupport@stemcell.com.

Notes

ACTIVITY UNITS

Collagenase: 1 collagenase digestion unit (CDU) equals 1 μmol of L-leucine equivalents released from collagen in 5 hours at 37°C, pH 7.5.

Caseinase: 1 unit equals 1 μmol of L-leucine equivalents released from 25 mg vitamin-free casein in 5 hours at 37°C, pH 7.5. Measures non-specific proteolytic activity.

Clostripain: 1 unit hydrolyzes 1 μmol of N α -benzoyl-L-arginine ethyl ester (BAEE)/minute at 25°C at pH 7.6, after activation in 2.5 mM dithiothreitol (DTT).

Trypsin: 1 unit hydrolyzes 1 μmol of BAEE/minute at 25°C at pH 7.6.

References

- Barthel ER et al. (2012) Tissue engineering of the intestine in a murine model. *J Vis Exp* (70): e4279.
- Comhair SAA et al. (2012) Human primary lung endothelial cells in culture. *Am J Respir Cell Mol Biol* 46(6): 723–30.
- Dong G et al. (2015) FOXO1 Regulates Dendritic Cell Activity through ICAM-1 and CCR7. *J Immunol* 194(8): 3745–55.
- Kessler E & Yaron A. (1973) A novel aminopeptidase from clostridium histolyticum. *Biochem Biophys Res Commun* 50(2): 405–12.
- Lee B et al. (2014) Anoctamin 1 contributes to inflammatory and nerve-injury induced hypersensitivity. *Mol Pain* 10(1): 5.
- Mohapatra A et al. (2016) Group 2 innate lymphoid cells utilize the IRF4-IL-9 module to coordinate epithelial cell maintenance of lung homeostasis. *Mucosal Immunol* 9(1): 275–86.
- Satish L et al. (2015) Expression analysis of human adipose-derived stem cells during in vitro differentiation to an adipocyte lineage. *BMC Med Genomics* 8: 41.
- Sharon Y et al. (2013) Isolation of normal and cancer-associated fibroblasts from fresh tissues by fluorescence activated cell sorting (FACS). *J Vis Exp* (71): e4425.
- Wang J et al. (2015) Interleukin 18 function in atherosclerosis is mediated by the interleukin 18 receptor and the Na-Cl co-transporter. *Nat Med* 21(7): 820–6.
- Wu Z et al. (2015) Depletion of MEIS2 inhibits osteogenic differentiation potential of human dental stem cells. *Int J Clin Exp Med* 8(5): 7220–30.
- You Y & Brody SL. (2013) Culture and differentiation of mouse tracheal epithelial cells. *Methods Mol Biol* 945: 123–43.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.