

Dissociation Reagents

Elastase

For digestion of native elastin

Catalog # 07453
07454

25 mg
100 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Elastase is a serine protease consisting of a single polypeptide chain of 240 amino acid residues and contains four disulfide bridges (Shotton & Hartley). Elastase is synthesized as a zymogen (proelastase) and is converted to the active form by limited trypsin proteolysis at its N-terminal. Elastase preferentially cleaves the peptide bond of C-terminal neutral, non-aromatic amino acid residues (Schellenberger et al.). Elastase is frequently used in combination with other proteases such as collagenase and trypsin to digest native elastin.

Product Information

Alternative Names:	Elastoproteinase; Lysosomal elastase; Pancreatic elastase I; Pancreatopeptidase E
Format:	Lyophilized powder
Storage:	Store at 2 - 8°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Reconstitution:	Dissociation reagents can be reconstituted in a balanced salt solution or buffer of choice.
Molecular Weight:	26.0 kDa
CAS Number:	39445-21-1
Optimum pH:	8.5
Cleavage Site:	-X-†-Y- : X = uncharged, non-aromatic Y = nonspecific

Specifications

Source:	Porcine pancreas
Activity:	≥ 3 units/mg protein. See Notes for further information.

Related Products

For a complete list of dissociation reagents, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com or contact us at techsupport@stemcell.com.

Notes

ACTIVITY UNITS

1 unit cleaves 1 μ mol of N-succinyl-L-alanyl-L-alanyl-L-alanine-p-nitroanilide/minute at 25°C at pH 8.0.

References

- Barkauskas CE et al. (2013) Type 2 alveolar cells are stem cells in adult lung. *J Clin Invest* 123(7): 3025–36.
- Grinnell F & Zhu M. (1994) Identification of neutrophil elastase as the proteinase in burn wound fluid responsible for degradation of fibronectin. *J Invest Dermatol* 103(2): 155–61.
- Klein D et al. (2011) Vascular wall-resident CD44+ multipotent stem cells give rise to pericytes and smooth muscle cells and contribute to new vessel maturation. *PLoS One* 6(5).
- Miri AK et al. (2015) Study of extracellular matrix in vocal fold biomechanics using a two-phase model. *Biomech Model Mechanobiol* 14(1): 49–57.
- Rock JR et al. (2011) Multiple stromal populations contribute to pulmonary fibrosis without evidence for epithelial to mesenchymal transition. *Proc Natl Acad Sci USA* 108(52): E1475–83.
- Schellenberger V et al. (1989) Characterization of the S¹-subsite specificity of porcine pancreatic elastase. *Eur J Biochem* 179(1): 161–3.
- Shotton DM & Hartley BS. (1973) Evidence for the amino acid sequence of porcine pancreatic elastase. *Biochem J* 131(4): 643–75.
- Takahashi A et al. (2014) Proteoglycans maintain lung stability in an elastase-treated mouse model of emphysema. *Am J Respir Cell Mol Biol* 51(1): 26–33.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.