

# Cytokines

## Mouse Recombinant SDF-1 alpha (CXCL12)

Stromal cell-derived factor 1 alpha

Catalog # 78121  
78121.1

5 µg  
25 µg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## Product Description

Stromal cell-derived factor 1 alpha (SDF-1 $\alpha$ ) is a member of the CXC group of chemokines that binds to the G-protein coupled receptor, CXCR4, to regulate migration, proliferation, differentiation, and survival of many cell types including hematopoietic stem cells, B cells, and T cells. It is produced by bone marrow stromal cells, osteoblasts, endothelial cells, and neuronal cells. SDF-1 $\alpha$  was first identified as the pre-B-cell growth-stimulating factor (PBSF) in the mouse bone marrow-derived stromal cell line, PA6, in the growth of B cell precursors (Hayashi et al.). SDF-1 $\alpha$  primarily regulates cell motility during development and adulthood, including the homing of hematopoietic stem cells and neutrophils to fetal bone marrow during ontogeny (Ara et al. 2003a) and the recruitment of endothelial progenitor cells from bone marrow during angiogenesis in adulthood (Zheng et al.). In addition to its role in hematopoiesis, the SDF-1 $\alpha$ /CXCR4 signaling pathway is also essential for the homing of primordial germ cells to gonads (Ara et al. 2003b), the migration of granule cells in the cerebellum during neurogenesis (Zou et al.), and the migration of breast cancer cells to sites of metastasis (Muller et al.).

## Product Information

**Alternative Names:** CXCL12, PBSF, SDF-1  $\alpha$ , Stromal cell-derived factor-1  
**Accession Number:** Q4FJL5  
**Amino Acid Sequence:** KPVLSYSYC PCRFFESHIA RANVKHLKIL NTPNCALQIV ARLKNNNRQV CIDPKLKIWIQ EYLEKALNK  
**Predicted Molecular Mass:** 8 kDa  
**Species:** Mouse  
**Cross Reactivity:** Human  
**Formulation:** Lyophilized after dialysis against phosphate-buffered saline.  
**Source:** CHO


## Specifications

**Activity:** The specific activity is  $\geq 6.7 \times 10^2$  units/mg ( $EC_{50} \leq 1.5 \mu\text{g/mL}$ ) as determined by  $Ca^{2+}$  mobilization assay using CHO-K1 cells stably expressing human G $\alpha$ 15 and mouse CXCR4.  
**Purity:**  $\geq 95\%$   
**Endotoxin Level:** Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is  $\leq 0.2$  EU/ $\mu\text{g}$  protein.

## Preparation and Storage

**Storage:** Store at  $-80^{\circ}\text{C}$ .  
**Stability:** Stable as supplied for 12 months from date of receipt.  
**Preparation:** Centrifuge vial before opening. Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. As a general guide, do not store at  $2 - 8^{\circ}\text{C}$  for more than 1 week or at  $-20^{\circ}\text{C}$  for more than 3 months. Avoid repeated freeze-thaw cycles.

## Data



(A) The biological activity of Mouse Recombinant SDF-1 alpha (CXCL12) was tested by its ability to mobilize  $\text{Ca}^{2+}$  in CHO-K1/ $\text{G}\alpha 15/\text{hCXCR4}$  cells (human  $\text{G}\alpha 15$  and mCXCR4 stably expressed in CHO-K1 cells).  $\text{Ca}^{2+}$  mobilization was measured using a fluorometric assay method. The  $\text{EC}_{50}$  is defined as the effective concentration of the growth factor at which  $\text{Ca}^{2+}$  mobilization is at 50% of maximum. The  $\text{EC}_{50}$  in the above example is less than  $1.5 \mu\text{g/mL}$ .

(B)  $2 \mu\text{g}$  of Mouse Recombinant SDF-1 alpha (CXCL12) was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Mouse Recombinant SDF-1 alpha (CXCL12) has a predicted molecular mass of 8 kDa.

## Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit [www.stemcell.com/cytokines](http://www.stemcell.com/cytokines) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## References

- Ara T et al. (2003a) Long-term hematopoietic stem cells require stromal cell-derived factor-1 for colonizing bone marrow during ontogeny. *Immunity* 19: 257–67.
- Ara T et al. (2003b) Impaired colonization of the gonads by primordial germ cells in mice lacking a chemokine, stromal cell-derived factor-1 (SDF-1). *Proc Natl Acad Sci* 100(9): 5319–23.
- Hayashi S et al. (1990) Stepwise progression of B lineage differentiation supported by interleukin 7 and other stromal cell molecules. *J Exp Med* 171(5): 1683–95.
- Muller A et al. (2001) Involvement of chemokine receptors in breast cancer metastasis. *Nature* 410: 50–56.
- Zheng H et al. (2007) Migration of endothelial progenitor cells mediated by stromal cell-derived factor-1alpha/CXCR4 via PI3K/Akt/eNOS signal transduction pathway. *J Cardiovasc Pharmacol*. 50(3): 274–80.
- Zou YR et al. (1998) Function of the chemokine receptor CXCR4 in haematopoiesis and in cerebellar development. *Nature* 393: 591–4.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.